

# **EXHIBIT 17**

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
1 August 2002 (01.08.2002)

PCT

(10) International Publication Number  
**WO 02/058711 A1**

(51) International Patent Classification<sup>7</sup>: **A61K 31/715**

(21) International Application Number: PCT/US02/01276

(22) International Filing Date: 15 January 2002 (15.01.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/261,911 16 January 2001 (16.01.2001) US

(63) Related by continuation (CON) or continuation-in-part (CIP) to earlier application:  
US 60/261,911 (CIP)  
Filed on 16 January 2001 (16.01.2001)

(71) Applicant (for all designated States except US):  
**SLOAN-KETTERING INSTITUTE FOR CANCER RESEARCH** [US/US]; 1275 York Avenue, New York, NY 10021 (US).

(72) Inventor; and

(75) Inventor/Applicant (for US only): **CHEUNG, Nai-Kong, V.** [US/US]; 3 Glen Park Road, Purchase, NY 10577 (US).

(74) Agents: **CHAN, Albert, Wai-Kit** et al.; Law Offices of Albert Wai-Kit Chan, LLC, World Plaza, Suite 604, 141-07 20th Avenue, Whitestone, NY 11357 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),

European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

#### Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)
- of inventorship (Rule 4.17(iv)) for US only

#### Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: THERAPY-ENHANCING GLUCAN

(57) Abstract: This invention provides a composition comprising an effective amount of glucan capable of enhancing efficacy of antibodies. This invention further provides the above compositions and a pharmaceutically acceptable carrier. This invention also provides a method for treating a subject with cancer comprising administering the above-described composition comprising effective amount of glucan capable of enhancing efficacy of vaccines. This invention provides a composition comprising effective amount of glucan capable of enhancing efficacy of vaccines. This invention also provides a method of treating a subject comprising administering the above pharmaceutical composition to the subject. This invention provides a composition comprising effective amount of glucan capable of enhancing efficacy of natural antibodies. This invention provides a composition comprising effective amount of glucan capable of enhancing host immunity. This invention also provides a composition comprising effective amount of glucan capable of enhancing the action of an agent in preventing tissue rejection.



WO 02/058711 A1

**THERAPY-ENHANCING GLUCAN**

Throughout this application, various references are referred to. Disclosures of these publications in their entireties are hereby incorporated by reference into this application to more fully describe the state of the art to which this invention pertains.

**BACKGROUND OF THE INVENTION**

Monoclonal antibodies (MoAb) selective for tumors have therapeutic potential.<sup>1,2</sup> The introduction of hybridoma technology by Kohler and Milstein in 1975<sup>3</sup> and the advances in molecular biologic techniques have greatly expanded the potential of MoAb in human cancers. Anti-CEA antibody in colorectal cancer,<sup>4</sup> anti-CD20 antibodies in lymphoma,<sup>5</sup> anti-HER2 antibodies in breast cancer,<sup>6</sup> anti-tenascin antibodies in glial brain tumors,<sup>7</sup> MoAb M195 against CD33 in acute leukemia<sup>8</sup> and anti-TAG-72 antibodies in colon cancer<sup>9</sup> have shown efficacy in clinical trials. Our laboratory has developed the MoAb 3F8 which targets the ganglioside GD2 overexpressed on neuroblastoma. 3F8 has been shown to have a high specificity and sensitivity in the radioimmunodetection of minimal residual disease (MRD) in patients with NB,<sup>10</sup> and a significant impact when used as adjuvant therapy.<sup>11</sup>

The immune basis of clinical tumor response to MoAb is at least two fold, direct cytotoxicity and induced immunity. Antibody dependent cell-mediated cytotoxicity (ADCC) and complement-mediated cytotoxicity (CMC) are responsible for the direct killing of tumor cells. On the other hand, through tumor opsonization<sup>12</sup> or idiotype network,<sup>13</sup> tumor-specific immunity is induced. With this paradigm, how the body eliminates microbial pathogens remains highly relevant in our strategic approach to cancer therapy. Since the first description of innate immunity and acquired immunity model, several components have emerged center stage.<sup>14</sup>

Antibodies, complement, phagocytes, and "danger" receptors are core elements of innate immunity while antigen-presenting cells, T and B lymphocytes constitute essential players in acquired immunity. Despite the availability of tumor-selective monoclonal antibodies and the ample supply of phagocytes/natural killers, shrinkage of established tumors following antibody treatment alone, and the acquisition of specific immunity, are not common in both preclinical models and cancer patients. The absence of a danger signal and the diminution of complement action by complement resistance proteins on tumor cells may explain the inefficiency of antibody mediated clinical responses.<sup>15</sup> LPS and beta-glucan, being cell wall components of bacteria and fungus, respectively, are potent danger signals to the immune systems in all life-forms, from *Drosophila* to man.<sup>16</sup> While LPS is too toxic for human use,  $\beta$ -glucan is a relatively benign structural component extractable from cereals, mushrooms, seaweed and yeasts.<sup>17</sup> They are made up of 1,3- $\beta$ -D-glucopyranosyl residues along which are randomly dispersed single  $\beta$ -D-glucopyranosyl units attached by 1,6-linkages, giving a comb-like structure. The 1,3- $\beta$ -backbone and the 1,6-linked branches were thought to be important for their immune effects. Lentinan (from *Lentinus edodes*, Basidiomycete family) with 1,6 branches at mean of 3 main chain units, is licensed Japan for cancer treatment. Schizophyllan (from *Schizophyllum commune*, Basidiomycete family) and  $\beta$ -glucan from Baker's yeast (*Saccharomyces cerevisiae*) have also similar structures. From seaweed, Laminarin (1,3  $\beta$ -D-glucan with 1,6- $\beta$  side chain branching on every 10 glucose subunit along the polymer backbone) has been extracted. Because of its smaller size and water solubility, it was thought to have potential biologic utility. On the other hand  $\beta$ -glucan from barley, oat or wheat have mixed 1,3- $\beta$  and 1,4- $\beta$ -linkage in the backbone,

but no 1,6- $\beta$  branches, and generally higher molecular weights and viscosities. In addition, they have not yet been tested for their in vivo immunomodulatory effects in cancer models.

5 We now report our findings that oral beta-glucans derived from barley or oats can greatly enhance the anti-tumor activity of anti-tumor monoclonal antibodies in xenograft models. Given the low toxicity of oral  $\beta$ -glucan, their role  
10 in cancer therapy deserves careful study.

**SUMMARY OF THE INVENTION**

This invention provides a composition comprising an effective amount of glucan capable of enhancing efficacy of antibodies. In an embodiment, the antibody is a monoclonal antibody. In a further embodiment, the antibody is an antibody against cancer.

The cancer is recognized by antibodies, and which includes but not limited to neuroblastoma, melanoma, non-Hodgkin's lymphoma, Epstein-Barr related lymphoma, Hodgkin's lymphoma, retinoblastoma, small cell lung cancer, brain tumors, leukemia, epidermoid carcinoma, prostate cancer, renal cell carcinoma, transitional cell carcinoma, breast cancer, ovarian cancer, lung cancer colon cancer, liver cancer, stomach cancer, and other gastrointestinal cancers.

This invention further provides the above compositions and a pharmaceutically acceptable carrier, thereby forming pharmaceutical compositions.

This invention also provides a method for treating a subject with cancer comprising administering the above-described composition to the subject.

This invention provides a composition comprising effective amount of glucan capable of enhancing efficacy of vaccines. In an embodiment, the vaccine is against cancer. This invention also provides the above compositions and a pharmaceutically acceptable carrier, thereby forming a pharmaceutical composition.

This invention also provides a method of treating a subject comprising administering the above pharmaceutical composition to the subject. In an embodiment, the subject is a human subject. In an embodiment, the vaccine is against

infectious agents. The infectious agents include but are not limited to bacteria, viruses, fungi, or parasites.

5 This invention provides a composition comprising effective amount of glucan capable of enhancing efficacy of natural antibodies. In an embodiment, the antibodies are against cancer. In another embodiment, the antibodies are against infectious agents. The infectious agents include but are not limited to bacteria, viruses, fungi, or parasites.

10 This invention provides a composition comprising effective amount of glucan capable of enhancing host immunity. In another embodiment, the immunity is against cancer or infectious agents.

15 This invention also provides a composition comprising effective amount of glucan capable of enhancing the action of an agent in preventing tissue rejection. In another embodiment, the tissue is transplanted tissue or  
20 transplanted organ. In another embodiment, the tissue is the host as in graft-versus-host reactions.

This invention also provides the above compositions, wherein the glucan are. 1,3-1,4 mixed linkage, without 1,6 branches.

25 The invention further provides the above compositions, wherein the glucan is of high molecular weight. In an embodiment, the molecular weight of the glucan ranges from 250,000 to 450,000 daltons. This invention provides the  
30 above compositions, wherein the glucan is derived from barley, oat, wheat or moss.

This invention provides the above compositions, wherein the glucan is stable to heat treatment. In an embodiment, the  
35 composition is stable after boiling for 3 hours.

This invention provides the above compositions, wherein oral route is adopted when taken into a subject. In an embodiment, the effective dose is about  $\geq 25$  mg/kg/day, five days a week for a total of 2-4 weeks.



**DETAILED DESCRIPTION OF THE FIGURES****First Series Of Experiments**

5 **Figure 1. Synergistic effect of MoAb and  $\beta$ -glucan in LAN-1**  
Two million LAN-1 neuroblastoma cells were xenografted  
subcutaneously in Balb/c athymic mice. Treatment started in  
groups of 5 mice each, 2 weeks after tumor implantation when  
visible tumors reached 0.7-0.8 cm diameter. 3F8 group  
10 (solid circles) was treated with 200 ug of intravenous 3F8  
injected through the retroorbital plexus twice weekly (M and  
Th). 3F8 + BG group (open circle) was treated with 200  
ug i.v. 3F8 twice weekly plus oral Barley  $\beta$ -glucan (BG medium  
viscosity) 400 ug daily by gavage for a total of 21 days.  
15 BG group (open triangle) received  $\beta$ -glucan alone, 400 ug po  
daily for 21 days. Tumor size was measured from the first  
day of treatment, and the product of the largest diameters  
expressed as percent of that on day 0 of treatment. While  
BG alone and 3F8 alone showed no anti-tumor effect, the  
20 BG+3F8 group showed highly significant tumor shrinkage and  
suppression ( $p < 0.001$ ).

**Figure 2. Synergistic effect of MoAb and  $\beta$ -glucan in NMB-7**  
Experiment in figure 1 was repeated with the neuroblastoma  
25 cell line NMB7, a slower growing line. Again BG alone (open  
triangle) and 3F8 alone (solid circles) showed no anti-tumor  
effect, the BG+3F8 group (open circle) showed highly  
significant tumor shrinkage and suppression ( $p < 0.001$ ). Y-  
axis is relative tumor size in percent and X-axis the number  
30 of days from first treatment.

**Figure 3. Dose response of intraperitoneal (ip)  $\beta$ -glucan**  
Two million NMB-7 xenografted nude mice were treated at the  
time of visible tumors with 3F8 alone, normal saline  
35 control, or 3F8 plus increasing doses of intraperitoneal BG  
(4 ug [solid diamond], 40 ug [open square], 400 ug [large

open circle)) or 400 ug of po BG [small open circle], or 400 ug of ip Lentinan [open diamond]. Highly significant tumor shrinkage and suppression was shown in the combination groups except at 4 ug of BG dose. Oral BG appeared to be  
5 more effective than ip BG.

**Figure 4. Dose response of oral  $\beta$ -glucan** NMB-7 xenografted in nude mice were treated as in figure 3 except that dose response of oral  $\beta$ -glucan (4ug [open diamond], 40 ug [open  
10 triangle], 400 ug [open circle]) was compared to 400 ug of ip BG [solid square]. Control group received saline [solid circle]. 400 ug po was again highly significant in eradicating or suppressing tumor growth. 400 ip appeared to be as effective as 40 ug po. 4 ug was the least effective.

15 **Figure 5. Dose response of oral  $\beta$ -glucan in LAN-1** Five million LAN-1 cells were planted subcutaneously. Tumor growth was more rapid compared to 2 million NMB-7 cells. Again 4 ug [solid squares], 40 ug [solid triangle] were no  
20 different from controls. Only 400 ug po [open circle] and 4000 ug po [open square] showed significant tumor eradication or suppression.

**Figure 6. Comparison of various  $\beta$ -glucans**  $\beta$ -glucan [400 ug po qd] derived from barley [7 days/wk open circle, M-F/week open triangle], Maitake mushrooms [solid triangles], laminarin [open squares] were compared in their synergism  
25 with antibody 3F8 against NMB-7 subcutaneous xenografts.

30 **Figure 7. More comparison of various  $\beta$ -glucans**  $\beta$ -glucans (400 ug po qd) from different barley lots [large open circle, small open circle], lentinan [open diamond], PSK [cross] were compared to mannan [solid square], 3F8 only [open triangle] or no treatment [solid circle]. Only BG

from barley showed synergistic anti-tumor effect with antibody 3F8 against LAN-1 xenografts.

**Figure 8. D-fraction Maitake Mushroom  $\beta$ -glucan** [open square] had no anti-tumor effect when compared to barley  $\beta$ -glucan alone [solid circle], 3F8 alone [open triangle], in contrast to barley  $\beta$ -glucan plus 3F8 [open circle] which was highly effective.

**Figure 9. Barley  $\beta$ -glucan of large molecular weight is more effective**  $\beta$ -glucan of 40K [solid square], 123K [open triangle], 183K [open square], 254K [open diamond], and 359K [open circle] were tested at 40 ug po daily dose in combination with 3F8 against LAN-1 subcutaneous xenografts. The larger the size of the  $\beta$ -glucan, the more effective the synergistic effect.

**Figure 10.  $\beta$ -glucans of low molecular weight and low viscosity was ineffective**  $\beta$ -glucans of various viscosities [40 ug po qd] derived from barley and oats were tested in combination with 3F8 against LAN-1 subcutaneous xenografts. Barley medium viscosity [large open circle], barley high viscosity [open square], oat medium viscosity [small open circle, dotted line], and oat high viscosity [open square, dotted line] were all effective in shrinking and suppressing tumor growth, in contrast to low viscosity barley  $\beta$ -glucan [solid square].

**Figure 11. Removal of NK cells by anti-Asialo GM1 antiserum in LAN-1 xenograft** decreased but did not eliminate the anti-tumor effect of barley  $\beta$ -glucan plus 3F8.

**Figure 12.** Removal of NK cells by anti-Asialo GM1 antiserum in NMB-7 xenograft again decreased but did not eliminate the anti-tumor effect of barley  $\beta$ -glucan plus 3F8.

5 **Figure 13.** 3F8-F(ab')<sub>2</sub> fragment [solid square], nonspecific human IgG [small solid square] or IgM [solid diamond] have no anti-tumor effect while 3G6 (IgM anti-GD2, open circle) was almost as effective as 3F8 (IgG3 anti-GD2, open triangle).

10 **Figure 14.** Barley  $\beta$ -glucan synergizes with R24 anti-GD3 antibody in SKMel28 melanoma xenografts in nude mice In contrast to  $\beta$ -glucan control [solid diamond], and R24 control [solid circle], the combination of R24 and  $\beta$ -glucan  
15 [open circle] significantly suppressed tumor growth.

**Figure 15.** Barley  $\beta$ -glucan synergizes with 3F8 anti-GD2 antibody against B16D14 murine melanoma in C57Bl/6 mice In contrast to saline control [solid circle],  $\beta$ -glucan control  
20 [solid triangle], and 3F8 control [solid square], the combination of 3F8 and  $\beta$ -glucan [open circle] significantly suppressed tumor growth.

**Figure 16.** Barley  $\beta$ -glucan synergizes with 3F8 anti-GD2  
25 antibody against B16D14-KbKd murine melanoma in C57Bl/6 mice In contrast to 3F8 control [solid circle], the combination of 3F8 and  $\beta$ -glucan [open circle] significantly suppressed tumor growth.

30 **Figure 17.** Barley  $\beta$ -glucan plus 3F8 did not affect GD2-negative B16 melanoma in C57Bl/6 mice The combination of 3F8 and  $\beta$ -glucan [solid circle] did not significantly suppress tumor growth when compared to controls [open circle].

**Figure 18. Barley  $\beta$ -glucan synergizes with 3F8 anti-GD2 antibody against E14 murine lymphoma in C57Bl/6 mice** In contrast to control [solid circle], the combination of 3F8 and  $\beta$ -glucan [open circle] significantly suppressed tumor growth.

**Figure 19. BARLEY Glucan syngergizes with 3F8 in prolonging survival from NMB7 neuroblastoma.** Nude mice (n=21) with established neuroblastoma NMB7 xenografts treated with 3F8 and barley  $\beta$ -glucan (open circles) had significantly longer median survival >300 days compared to 30 days in the control mice (solid triangle) treated with saline alone, 3F8 alone, or  $\beta$ -glucan alone (p<0.001). Long-term survival was 56% in the treatment group and 5% in the control group.

**Figure 20. BARLEY glucan synergizes with 3F8 in prolonging survival from LAN-1 neuroblastoma.** In nude mice bearing established LAN1 xenografts median survival increased from 20 days in the control group (n=38, solid triangles) to 42 days in the 3F8 plus glucan group (n=48, open circles, p<0.001).

### Second Series of Experiments

**Figure 1. Synergistic effect of MoAb and  $\beta$ -glucan in neuroblastoma xenografts.** Two million neuroblastoma cells (1A: LAN-1, 1B: NMB7, 1C:SK-N-ER) were xenografted subcutaneously in athymic Balb/c mice. Treatment started in groups of 5 mice each, 2 weeks after tumor implantation when visible tumors reached 0.7-0.8 cm diameter. 3F8 group (solid circles) was treated with 200 ug of intravenous 3F8 injected through the retroorbital plexus twice weekly (M and Th). 3F8 + BG group (open circle) was treated with 200 ugi.v.3F8 twice weekly plus oral  $\beta$ -glucan (BG) 400 ug daily by gavage for a total of 21-29 days. BG group (open triangle) received 400 ug oral  $\beta$ -glucan alone. Tumor size

was measured from the first day of treatment, and the product of the largest diameters expressed as percent of the size on day 0 of treatment. Vertical bars represent standard errors, which were similar for the glucan and the 3F8 alone groups. While BG alone and 3F8 alone showed no anti-tumor effect, the BG+3F8 group showed highly significant tumor shrinkage and suppression ( $p < 0.01$ ).

**Figure 2. Dose response of intraperitoneal (ip)  $\beta$ -glucan.**

Two million NMB7 xenografted athymic nude mice were treated at the time of visible tumors with 3F8 alone, normal saline control, or 3F8 plus increasing doses of intraperitoneal BG (4 ug [solid diamond], 40 ug [open square], 400 ug [large open circle]) or 400 ug of po BG [small open circle]. Highly significant tumor shrinkage and suppression was shown in the combination groups except at 4 ug of BG dose. Oral BG appeared to be more effective than ip BG.

**Figure 3. Dose response of oral  $\beta$ -glucan.** NMB7 xenografted in nude mice were treated as in Figure 3 except that dose response of oral  $\beta$ -glucan (4ug [open diamond], 40 ug [open triangle], 400 ug [open circle]) was compared to 400 ug of ip BG [solid square]. Control group received saline [solid circle]. 400 ug po was again highly significant in eradicating or suppressing tumor growth. 400 ip appeared to be as effective as 40 ug po. 4 ug was the least effective.

**Figure 4. Removal of NK cells by anti-Asialo GM1 antiserum on  $\beta$ -glucan effect in LAN-1 xenografts** decreased but did not eliminate the anti-tumor effect of  $\beta$ -glucan plus 3F8.

**Figure 5. Barley  $\beta$ -glucan synergized with 3F8 in prolonging survival from NMB7 neuroblastoma.** Nude mice (n=22) with established neuroblastoma NMB7 xenografts treated with 3F8 and barley  $\beta$ -glucan (solid line) had significantly longer median survival (median 166 days)

compared to control mice (n=34, broken line, median 30 days) treated with saline alone (n=10), 3F8 alone (n=8), or  $\beta$ -glucan alone (n=16) ( $p<0.001$ ). Long-term survival was 47% in the treatment group and 3% in the control group.

**Figure 6. Barley  $\beta$ -glucan synergized with 3F8 in prolonging survival from LAN-1 neuroblastoma.** In nude mice bearing established LAN-1 xenografts, median survival increased from 21 days in the control group (n=55, broken line) to 54 days in the 3F8 plus glucan group (n=82, solid line,  $p<0.001$ ).

### Third Series of Experiments

**Figure 1. Oral  $\beta$ -glucan synergizes with 3F8 in prolonging survival from neuroblastoma.** Nude mice (n=22, solid line) with established NMB7 xenografts (0.7-0.8 cm diameter tumor at the beginning of treatment) were treated with 3F8 (200 ug twice a week iv) and 400 ug of  $\beta$ -glucan po daily for a total of 3 weeks. Control mice received either saline alone (n=10, broken line), 3F8 alone (n=8, dashed line), or  $\beta$ -glucan (n=16, dotted line) alone. Median survival was 30 days in control groups and 166 days in the treatment (3F8 plus  $\beta$ -glucan, n=22) group ( $p<0.001$ ). Ten (45%) in the combination group survived long term with a median follow-up of 248 days. Only one mouse in any of the control groups (<5%) remained alive during the experiment.

**Figure 2. Synergy of Oral barley  $\beta$ -glucan with (A) R24 (anti-GD3) antibody against SKMel28 melanoma xenografts in nude mice.** In contrast to  $\beta$ -glucan control [solid diamonds], and R24 control [solid circles], the combination of R24 and  $\beta$ -glucan [open circles] significantly suppressed tumor growth (tumor growth rate reduced for combination treatment by 1.2%, 95% CI -0.1%, 2.5%,  $p=0.06$ ) (B) 528 (anti-EGF-R) MoAb against epidermoid carcinoma A431 xenografts in nude mice. In contrast to  $\beta$ -glucan + 455

(IgG1 noncomplement fixing) control [solid squares], and 528 MoAb alone [solid circles], the combination of 528 MoAb and  $\beta$ -glucan [open circles] significantly suppressed tumor growth (tumor growth rate reduced for combination treatment by 1.4%, 95% CI -0.7%, 3.5%,  $p=0.17$ ). (C) Herceptin (anti-HER2) antibody against human breast carcinoma BT474 xenografts in nude mice. In contrast to control [n=4, solid circles], Herceptin [n=9, open squares], or  $\beta$ -glucan control [n=7, solid squares], the combination of Herceptin and  $\beta$ -glucan [n=12, open circles] significantly suppressed tumor growth (tumor growth rate reduced for combination treatment by 1.9%, 95% CI 0.7%, 3%,  $p=0.002$ ).



**DETAILED DESCRIPTION OF THE INVENTION**

This invention provides a composition comprising an effective amount of glucan capable of enhancing efficacy of antibodies.

In an embodiment, the antibody is a monoclonal antibody. In a further embodiment, the antibody is an antibody against cancer. In another embodiment, the antibody is a tumor-binding antibody. In a further embodiment, the antibody is capable of activating complement. In a still further embodiment, the antibody is further capable of activating the antibody dependent cell-mediated cytotoxicity.

In an embodiment, the antibody is directed at the epidermal growth factor receptor. In a further embodiment, the antibody is 528 or C225.

In another embodiment, the antibody is directed to a ganglioside. In a further embodiment, the ganglioside is GD3. In a still further embodiment, the antibody is R24.

In a separate embodiment, the ganglioside is GD2. In a further embodiment, the antibody is 3F8.

In an embodiment, the antigen recognized by the antibody is CD20. In a further embodiment, the antibody is Rituximab.

In another embodiment, the antigen is CD25. In a further embodiment, the antibody is Dacluzimab.

In a separate embodiment, the antigen is Her2/neu. In a further embodiment, the antibody is Herceptin.

In another embodiment, the antigen is CD22. In a further embodiment, the antibody is Epratuzumab.

The cancer is recognized by antibodies, and which includes but not limited to neuroblastoma, melanoma, non-Hodgkin's lymphoma, Epstein-Barr related lymphoma, Hodgkin's lymphoma, retinoblastoma, small cell lung cancer, brain tumors, leukemia, epidermoid carcinoma,<sup>40</sup> prostate cancer,<sup>40,41</sup> renal cell carcinoma,<sup>40</sup> transitional cell carcinoma,<sup>40</sup> breast cancer,<sup>42,43</sup> ovarian cancer,<sup>40</sup> lung cancer, colon cancer,<sup>40</sup> liver cancer, stomach cancer, and other gastrointestinal cancers.

10 This invention further provides the above compositions and a pharmaceutically acceptable carrier, thereby forming pharmaceutical compositions.

15 This invention also provides a pharmaceutical composition comprising a combination as described above and a pharmaceutically acceptable carrier. For the purposes of this invention, "pharmaceutically acceptable carriers" means any of the standard pharmaceutical carriers. Examples of  
20 suitable carriers are well known in the art and may include, but are not limited to, any of the standard pharmaceutical carriers such as a phosphate buffered saline solution and various wetting agents. Other carriers may include additives used in tablets, granules and capsules, etc. Typically such  
25 carriers contain excipients such as starch, milk, sugar, certain types of clay, gelatin, stearic acid or salts thereof, magnesium or calcium stearate, talc, vegetable fats or oils, gum, glycols or other known excipients. Such carriers may also include flavor and color additives or  
30 other ingredients. Compositions comprising such carriers are formulated by well-known conventional methods.

This invention also provides a method for treating a subject with cancer comprising administering the above-described  
35 composition to the subject.

This invention provides a composition comprising effective amount of glucan capable of enhancing efficacy of vaccines. In an embodiment, the vaccine is against cancer.

5 This invention also provides the above compositions and a pharmaceutically acceptable carrier, thereby forming a pharmaceutical composition.

10 This invention also provides a method of treating a subject comprising administering the above pharmaceutical composition to the subject. In an embodiment, the subject is a human subject.

In an embodiment, the vaccine is against infectious agents.  
15 The infectious agents include but are not limited to bacteria, viruses, fungi, or parasites.

This invention provides a composition comprising effective amount of glucan capable of enhancing efficacy of natural  
20 antibodies.

In an embodiment, the antibodies are against cancer.

In another embodiment, the antibodies are against infectious  
25 agents. The infectious agents include but are not limited to bacteria, viruses, fungi, or parasites.

This invention provides a composition comprising effective amount of glucan capable of enhancing host immunity. In  
30 another embodiment, the immunity is against cancer or infectious agents.

This invention also provides a composition comprising effective amount of glucan capable of enhancing the action  
35 of an agent in preventing tissue rejection.

In an embodiment, the agent is an antibody. In a further embodiment, the antibody modulates T-cell function. In a

still further embodiment, the antibody is anti-CD25 or anti-CD3.

5 In a separate embodiment, the antibody modulates B-cell function. In another embodiment, the antibody is anti-CD20.

In another embodiment, the tissue is transplanted tissue or transplanted organ. In another embodiment, the tissue is the host as in graft-versus-host reactions.

10 This invention also provides the above compositions, wherein the glucan are. 1,3-1,4 mixed linkage, without 1,6 branches.

15 The invention further provides the above compositions, wherein the glucan is of high molecular weight. In an embodiment, the molecular weight of the glucan ranges from 250,000 to 450,000 daltons.

20 This invention provides the above compositions, wherein the glucan is derived from barley, oat, wheat or moss.

This invention provides the above compositions, wherein the glucan is stable to heat treatment. In an embodiment, the composition is stable after boiling for 3 hours.

25 This invention provides the above compositions, wherein oral route is adopted when taken into a subject. In an embodiment, the effective dose is about  $\geq 25$  mg/kg/day, five days a week for a total of 2-4 weeks.

30 The invention will be better understood by reference to the Experimental Details which follow, but those skilled in the art will readily appreciate that the specific experiments detailed are only illustrative, and are not meant to limit  
35 the invention as described herein, which is defined by the claims which follow thereafter.

## EXPERIMENTAL DETAILS

**Materials And Methods**

**Cell lines** Human neuroblastoma cell lines LA-N-1 was provided by Dr. Robert Seeger, Children's Hospital of Los Angeles, Los Angeles, CA and NMB7 by Dr. Shuen-Kuei Liao (McMaster University, Ontario, Canada). Neuroblastoma cell lines SKNHM, SKNHB, SKNJD, SKNLP, SKNER, SKNMM, SKNCH and SKNSH were derived from patients with metastatic disease treated at Memorial Sloan-Kettering Cancer Center (MSKCC), New York, NY. Other cells lines, Daudi, RPMI 6666, SKMel-28, A431, B16 were derived from American Type Culture Collection (ATCC), Rockville, MD. The cell lines B16, B16D14 were kindly provided by Dr. Kenneth Lloyd of Memorial Sloan-Kettering Cancer Center. Kb transfected (B16D14Kb) and (Kb + Kd) transfected (B16D14KbKd) were kindly provided by Dr. Michel Sadelain, MSKCC. Cell lines were cultured in 10% defined calf serum (Hyclone, Logan, UT) in RPMI with 2 mM L-glutamine, 100 U/ml of penicillin (Sigma, St. Louis, MO), 100 ug/ml of streptomycin (Sigma), 5% CO<sub>2</sub> in a 37°C humidified incubator. Normal human mononuclear cells were prepared from heparinized bone marrow samples by centrifugation across a Ficoll-Hypaque density separation gradient.

**Antibodies** Monoclonal antibodies 3F8 (mouse IgG3) and 3G6 (mouse IgM), and 8H9 (mouse IgG1) reactive with neuroblastoma have been previously described. They were produced by as ascites and purified by affinity chromatography: protein A (Pharmacia, Piscataway, NJ) for 3F8,<sup>18</sup> protein G (Pharmacia) for 8H9,<sup>19</sup> and Clq-sepharose (Pierce Chemicals) for 3G6.<sup>18,20</sup> These antibodies are >90% pure by SDS-PAGE. F(ab')<sub>2</sub> fragments were prepared by pepsin digestion as previously reported.<sup>21</sup> Anti-GD3 antibody (R24) was kindly provided by Dr. Paul Chapman of MSKCC.<sup>22</sup> FLOPC21, an IgG3 myeloma, was purchased from Sigma Chemicals, St. Louis, MI. TIB114 (N.S.7) a hybridoma

secreting an IgG3 control antibody was obtained from ATCC. Rabbit anti-asialo-GM1 antibody (Wako Pure Chemical Industries, Ltd, Osaka, Japan) diluted to 1 mg/ml of protein was administered at 200  $\mu$ l ip on days 0, 1, 2, 7, 14, 21.

5 Rituximab, anti-CD20 antibody was purchased from Genentech, Inc., CA.

**Indirect immunofluorescence** 1 million target cells were washed in PBS and then spun at 180 x g for 5 min. The pellets were then reacted with 100  $\mu$ l of 15 $\mu$ g/ml 8H9 at 4°C for 1 hour. After washing the cells with PBS they were allowed to react with 100 $\mu$ l FITC-conjugated goat F (ab<sup>2</sup>) anti- mouse IgG + IgM, (Biosource International, Camarillo, CA) at 4° C.<sup>18</sup> Flow cytometric analysis was performed using FACSCalibur Immunocytometer (Becton-Dickinson Immunocytometry Systems, San Jose, CA).

**Glucan** 1,3-1,4- $\beta$ -glucan derived from barley, 1,3-1,6-- $\beta$  glucan (Laminarin) from seaweed (Laminaria digitata), and mannan were purchased from Sigma Co. 1,3-1,6- $\beta$ -glucan (Lentinan) was obtained from Drug Synthesis and Chemistry Branch, Developmental Therapeutics Program, Division of Cancer Treatment, National Cancer Institute, Bethesda, MD. Maitake mushroom glucan (containing 1,3-1,6- $\beta$ -glucan extracted from Grifola frondosa) D-fraction was obtained from Maitake Products Inc., Paramus, NJ. Barley and oats  $\beta$ -glucans of various molecular sizes (measured by high performance size-exclusion chromatography [HPSEC] with multiple angle laser scattering [MALLS] detection) and viscosities (measured in cSt units) were obtained from Megazyme International Ireland Ltd., Bray, County, Ireland. Barley glucan was dissolved by boiling for 10 minutes in normal saline. A stock solution of Lentinan in DMSO (Sigma) was diluted in water before use.

**Mice and treatment** Athymic Balb/c and C57Bl/c mice were purchased from Jackson Laboratories, Bar Harbor, ME, and CB-

17 SCID from Taconic. Tumor cells were planted (1-5 x10<sup>6</sup>) cells) in 100ul of Matrigel (Sigma Co) subcutaneously. Following implantation, tumor sizes (maximum width and lengths) were measured. Tumor size was calculated as product of the 2 perpendicular diameters. Treatment studies started when tumor diameter reached 0.7 to 0.8 cm, usually by 14-21 days of tumor implantation. Mice received antibody treatment intravenously (by retroorbital injection) twice weekly and glucan by gavage every day for a total 3-4 weeks (21-8 days of glucan and 6-8 doses of antibody). Mice were weighed once a week and tumor size measured twice a week. Mice were sacrificed when tumors reached sizes that interfered with their well-being.

<sup>51</sup>Chromium (51Cr) release assay<sup>23</sup> In brief, 2x10<sup>3</sup> of 51Cr-labeled target cells were mixed with effector cells in a final volume of 0.2 ml of medium in 96-well flat-bottomed microtiter plates (Costar, Cambridge, MA). The plates were incubated for 4 h at 37°C in 5%CO<sub>2</sub> and then centrifuged. 100ul of assay supernatant was counted in a gamma counter. Target cell spontaneous chromium release ranged from 10 to 25%.

## RESULTS

Barley glucan synergizes with anti-GD2 antibody 3F8 in eradicating human neuroblastoma. 3F8 is a murine IgG3 monoclonal antibody specific for ganglioside GD2. It activates mouse and human complement, and mediates effective ADCC against human neuroblastoma cells in vitro. Barley glucan when administered orally at 400 ug qd had no appreciable effect on tumor growth compared to antibody 3F8 given i.v. alone. However, when barley glucan and 3F8 were used in combination, tumor growth was near totally suppressed. In >40% of mice, NMB7 tumors remained permanently suppressed even when treatment was stopped after 21 days. Similar observations were made with neuroblastoma

cell lines derived from different sources: NMB7, LAN-1 (Figures 1 and 2) and SK-N-ER (data not shown). Barley glucan was equally effective when administered orally or intraperitoneally. In contrast, for the GD2-negative  
5 rhabdomyosarcoma, HTB82, 3F8 plus  $\beta$ -glucan treatment was ineffective (data not shown).

**Dose response curve for ip barley glucan.** When the dose of intraperitoneal barley glucan was decreased 10-fold from 400  
10 ug, it was clear that 4 ug was no longer effective in synergizing with MoAb 3F8 in suppressing NMB7 growth. Interestingly, both ip lentinan and po glucan (at 400 ug po qd) were also effective (Figure 3).

**Oral barley glucan is as effective as ip glucan** When oral barley glucan was studied in NMB7 tumor (Figure 4) followed after treatment for 80 days, similar dose response was found, i.e. while 400 ug oral regimen was curative, breakthroughs were seen for the other dose levels, with 4 ug  
20 oral dose escaping sooner than 40 ug. Interestingly, 400 ug ip was only as effective as the 40 ug oral group, with late breakthrough tumor growths around the same time, unlike the 400 ug po group, where all tumors remained suppressed despite stopping all therapy after 21 days. Using the LAN1  
25 tumor model, both 4 and 40 ug glucan were ineffective compared to 400 and 4000 ug of glucan per dose (Figure 5). There was no significant body weight change in the treatment groups (after accounting for tumor growth) irrespective of dose of glucan or combination with antibody 3F8. At  
30 necropsy on day 21, there were no appreciable difference in the peripheral blood counts, cholesterol and blood chemistry between mice receiving different glucan doses. There was also no difference in the histologic appearances of organs in mice treated with glucan at any of the dose levels, when  
35 compared to control mice that received saline.



By the oral route, only certain glucans, and frequent dosing were effective. For NMB7 tumors (Figure 6), 400 ug oral maitake was effective in synergizing with antibody 3F8, although late breakthroughs were seen. A 5 day/week po barley glucan regimen was equally as effective as the daily regimen. In contrast, a once a week or twice a week schedule of barley glucan was ineffective (data not shown). For the faster growing LAN1 tumors (Figure 7) unlike barley glucan (lot #1 and lot #2), po lentinan, PSK or mannan were all ineffective. The effect of Maitake glucan was not significantly different from glucan dose or 3F8 alone (Figure 8). Glucan from barley was more effective than that from oat despite similarities in their molecular sizes (Figure 11).

**Barley glucans of large molecular weight is more effective**  
In Figure 9, barley glucans of different molecular sizes (40K, 123K, 183K, 254K, 359K) were tested at an oral dose of 40 ug. Anti-tumor effect improved with increasing molecular weights, such that glucan of 359K size was most effective. Nevertheless, at high doses (e.g. 400 ug) even the less effective sizes, 40K and 128K showed some benefit (092900megazyme.xls from folder megazyme). Glucan derived from oat also showed synergistic anti-tumor effect when administered in the presence of 3F8. Both molecular size and viscosity appeared to be important for this anti-tumor effect. For example barley glucan (327K, >100 cSt or 250K, 25 cSt) and oat glucan (69 cSt or 20-30 cSt) were highly effective in synergizing with MoAb, whereas barley glucan of 137K and 5.6 cSt was not (Figure 10).

**Role of NK cells in glucan effect.** Removal of NK cells using anti-Asialo GM1 antiserum eliminated a substantial amount, although not completely the anti-tumor activity of glucan (Figure 11 and 12). Moreover in beige mice glucan

was effective in synergizing with 3F8 (data not shown), suggesting that at least part of the anti-tumor activity was mediated by NK-independent cytotoxicity.

5 **IgG3-F(ab')<sub>2</sub> or IgG1 antibodies do not have substantial anti-tumor activity (Figure 13)** The role of Fc in mediating the anti-tumor effect of glucan was apparent when Fc was removed by pepsin or when IgG1 isotype (data not shown) was used. Neither was able to activate complement or mediate  
10 efficient ADCC, and neither has significant anti-tumor effect when administered with 400 ug of oral glucan.

**Barley Glucan synergizes with other complement fixing antibodies in a wide spectrum of human tumors.** IgG3 anti-  
15 GD3 antibody R24 synergized with po glucan in shrinking **melanoma SKMel-28** xenografts (Figure 14). Rituximab (humanized IgG1 anti-CD20) synergized with po glucan in eradicating **EBV-lymphoma, Daudi lymphoma, and Hodgkin's disease**. Although anti-EGF-R antibody 428 (mouse IgG2a)  
20 was able to suppress **epidermal carcinoma**<sup>40</sup> A431 tumor growth, 428 plus oral glucan was much more effective in eradicating tumors.

	<u>Cell line</u>	<u>Antibody</u>
25	<u>Subcutaneous tumor models</u>	
	<u><b>Human xenografts</b></u>	
	NMB7	3F8
	LAN-1	3F8
	SK-N-ER	3F8
	SK-N-	3F8
30	SKMel-28	R24
	EBV-lymphoma	Rituximab
	Daudi lymphoma	Rituximab
	Hodgkin's disease	Rituximab
	Epidermal Carcinoma 528	
35	<u><b>Syngeneic tumors</b></u>	
	EL4	3F8
	B16D14	3F8
	B16D14-Kb	3F8
40	B16D14-Kb-Kd	3F8

Metastatic tumor models

**Human xenografts**

Daudi

3F8

**Syngeneic tumors**

5

EL4

3F8

**Glucan synergizes with 3F8 in C57Bl/6 mice against syngeneic tumors.** While our early experiments were focused on human xenografts in athymic or SCID mice, similar synergism was observed in immunologically intact mice grafted with GD2-positive B16 melanomas (B16D14 [Figure 15], B16D14Kb, or B16D14KbKd [Figure 16]) or GD2-positive EL4 lymphoma [Figure 18]. Neither barley glucan nor 3F8 by itself showed anti-tumor effect. In contrast the combination of glucan and 3F8 was able to suppress almost entirely tumor growth in C57Bl/6 mice. In control B16 tumors which were GD2-negative, glucan plus 3F8 had no anti-tumor effect. We conclude that the glucan effect observed was not restricted to immune deficient animals. In addition, it requires tumor-specific antibodies, i.e. effective only if tumor cells carried the target antigen.

**Glucan synergizes with 3F8 in C57Bl/6 mice against metastatic tumors.** When EL4 lymphoma cells were injected iv, mice developed widespread tumors in their lungs, livers and at the site of injection and rapidly succumbed. Control animals were dead by 14 days following EL4 injection, while the group treated with i.v. 3F8 plus 400 ug glucan po (treatment initiated 5 days following EL4 injection) had significantly longer survival.

**Barley Glucan synergizes with 3F8 in prolonging survival** Nude mice (n=21) with established neuroblastoma NMB7 xenografts (0.7-0.8 cm diameter tumor at the beginning of treatment) were treated with 3F8 (200 ug twice a week iv) and 400 ug of barley  $\beta$ -glucan po daily for a total of 3 weeks. Control mice (n=21) received either saline alone,

3F8 alone, or  $\beta$ -glucan alone. Median survival was 30 days in control and >300 days in the treatment (3F8 plus  $\beta$ -glucan) group ( $p < 0.001$ ). Long-term survival was 56% in the treatment group and 5% in the control group (**Figure 19**).

5 In nude mice bearing established LAN1 xenografts (also 0.7-0.8 cm diameter tumor at the beginning of treatment) median survival increased from 20 days in the control group ( $n=38$ ) to 42 days in the 3F8 plus glucan group ( $n=48$ ) ( $p < 0.001$ , **Figure 20**).

10

### Discussion

Using the human xenograft and syngeneic mouse tumor models, we have made the following observations. Glucan derived from barley or oats can synergize with monoclonal antibodies  
15 in suppressing or eradicating tumors, while  $\beta$ -glucan or antibody alone has little anti-tumor effect. Anti-tumor response requires antibodies that activate complement, whether mouse IgM, mouse IgG3 or human IgG1. Glucans of high molecular weight 250K and viscosity (20 cSt) possess  
20 this special effect. Oral route is at least equally (if not more) effective than the intraperitoneal route. It is a dose-dependent phenomenon, where 400 ug per dose is required for maximal effect. Natural killer cells are not essential for this glucan phenomenon, although they contribute to the  
25 anti-tumor effect. Normal T-cells and B-cells are not required for the anti-tumor effect since immune-deficient mouse strains demonstrate the glucan effect, whether athymic, SCID or SCID-beige mice are used. In addition, normal T-cells and B-cells do not interfere with this glucan  
30 effect, as shown in the syngeneic C57Bl/6 mouse model. Most importantly, oral glucan is well-tolerated by all the mice tested so far, with no noticeable change in body weight, blood counts or organ histologies, even at doses as high as 4 mg per dose per day.

35

Our findings differ significantly from previous observations and predictions on the use of glucans in cancer treatment. In the past it was thought that the 1,3-1,6- $\beta$ linkage was absolutely required for the glucan anti-tumor effect.<sup>17</sup> This structure contains 1,3- $\beta$ -D-glucopyranosyl units along which are randomly dispersed single  $\beta$ -D-glucopyranosyl units attached by 1,6- linkages, giving a comb-like structure (e.g. Lentinan, Schizophyllan, Laminarin, and glucan from Baker's yeast). In these models, it was believed that T-cells cells were activated and indeed required for the anti-tumor effect. In addition, it was believed that small molecular weight glucan should be more effective than high molecular weight glucan and that the most effective administration should be intravenous or intraperitoneal routes. Indeed, Betafectin (PGG) was derived from a genetically engineered *Saccharomyces cerevisiae* which makes 1,3-1,6- $\beta$ -D glucans with weaker interchain associations.<sup>24</sup> It was manufactured for i.v. injection to improve macrophage function in the hope of reducing infectious complications and improving wound healing. Barley glucan is a linear polymer with 1,3 and 1,4 linkages; however, it is not a comb like structure. We did not find any anti-tumor effect of barley glucan when given alone. However, when used in combinations with monoclonal antibodies, the synergistic effect was remarkable. In addition, glucans of high molecular weight and high viscosity appeared to be most effective, contrary to what one might expect for macromolecular transport. Although barley glucan activates granulocyte mediated ADCC in vitro (data not shown), the effects of glucan may be indirect. It is not clear if the absorption of glucan is necessary for its anti-tumor effect. The exact mechanism of how barley glucan enhances the anti-tumor effect of monoclonal antibodies in vivo is unknown.

One possible mechanism of action may relate to innate receptors for  $\beta$ -glucan, in a hard-wired information network on phagocytes and lymphoid cells; receptors that normally recognize death signals and microbial molecular patterns.<sup>25</sup>

5 Monoclonal antibodies, either through Fc interaction or through CR3 interaction with iC3b, direct cytotoxicity to tumor cells, a process greatly enhanced by  $\beta$ -glucan activation of effector cells. This killing is immediate, nonclonal, and obligatory, a process often referred to as

10 innate immunity. The consequence of this innate effector arm is the activation of costimulatory molecules and induction of cytokines and chemokines that will enhance adaptive immunity to the tumor cells. Thus, activation of immunity is based upon discrimination between dangerous and

15 nondangerous antigens; and if cancer can be viewed as constant danger to the immune system,<sup>26,27</sup> memory T-cells will not become tolerized.  $\beta$ -glucan receptors belong to a family of pattern recognition receptors (PRRs) specific for pathogen-associated molecular patterns (PAMPs). They are

20 biosensors for invading pathogens widely distributed in vertebrate and invertebrate animals,<sup>28</sup> a nonclonal host defense pathway with structural and functional homologies in phylogenetic lineages that diverged over a billion years ago. A limited set of conserved signaling modules such as

25 Toll/IL-1R homology domain, the SIIK domain, the Rel homology domain and perhaps the leucine rich regions (LRR) domain, represent the original building blocks for PRRs. For example, insects respond to infection by rapid and transient synthesis of antimicrobial peptides by the fat

30 body and hemocytes. In drosophila antibacterial peptides (cecropin, attacin and defensin) and anti-fungal peptide drosomycin are dependent on the Toll pathway; this PRR activates a proteolytic cascade to act on Spatzle, or 18-Wheeler (18W) to form the active ligand for Toll.

Activation of the human Toll homologue results in induction of IL-1, IL-6, IL-8, B7.1 and B7.2. With B7, signaling through CD28 occurs; T-cell become activated followed by expression of surface molecules such as TNF- $\alpha$  and TNF- $\beta$ , Fas ligand (L), CD40L, CD30L, and CD27L, as well as secretion of cytokines. Interestingly, for dendritic cells, when they express B7, they stop antigen uptake (i.e. becoming nonendocytic) and assume their antigen-presenting role. Certain activation motifs such as LRR are present in Toll and the endotoxin receptor CD14; they are also present intracellularly in plants, probably responsible for resistance to intracellular pathogens.<sup>28</sup>

Carbohydrate-rich antigens on bacteria or fungi can activate complement. Alternatively, specific antibodies can also deposit complement components on pathogens or cancer cells, such as the C3b fragment of C3, which is rapidly proteolyzed into iC3b fragment by serum factor I. These iC3b fragments can glue pathogens or tumor cells to the iC3b-receptors (CR3, CD11b/CD18) on phagocytic cells and NK cells, stimulating phagocytosis and/or cytotoxic degranulation. Thus, antibody and complement link innate and adaptive immunity by targeting antigens to different cells of the immune system, e.g. via CR3 and Fc for phagocytic cells, CR2 for B cells, and CR1, CR2, or CR3 for follicular dendritic cells.<sup>29</sup> For neutrophils, CR3-dependent phagocytosis requires ligation of two distinct binding sites, one for iC3b and a second site for  $\beta$ -glucan. Without  $\beta$ -glucan, iC3b-opsonized target cells are resistant to killing.<sup>30</sup> Microbes possess polysaccharides that can activate the lectin domain on CR3, leading to phagocytosis and cytotoxic degranulation. In contrast, human cells (including tumors) lack these CR3-binding polysaccharides, thus the inability of CR3 to mediate phagocytosis or extracellular cytotoxicity of tumor cells opsonized with iC3b. The lectin site of CR3 can also

influence transmembrane signaling of endogenous neutrophil membrane GPI-anchored glycoproteins (CD14, CD16, CD59, CD87[uPAR]). In a mouse mammary tumor model, where there is naturally occurring IgM and IgG antibodies, injection of yeast soluble  $\beta$ -glucan could suppress tumor growth, an effect lost in C3-deficient or CD11b (CR3)-deficient mice.<sup>31,32</sup> Since iC3b bound to a primary protein antigen can also enhance recognition and specific antibody synthesis by antigen-specific B cells,<sup>33</sup> the presence of glucan plus complement activation may enhance B-cell response to pathogens or tumor cells.

If this synergistic effect of  $\beta$ -glucan on antibodies is active in humans, our findings may have broad clinical implications. First the efficacy of monoclonal antibodies in cancer (e.g. Herceptin, Rituximab, Dacluzimab, anti-GD2 and anti-EGF-R MoAb) can be potentially enhanced.<sup>34</sup> Nevertheless, even though toxicity from glucan is expected to be minimal, the enhanced efficacy of MoAb may also increase MoAb-mediated toxicity. For example, the side effects of Herceptin on cardiac function, or anti-GD2 MoAb on neuropathic pain may be increased. Second, since the amount and quality of barley and oat glucan in daily food intake can vary, future interpretations of efficacy trials using MoAb may need to take this into account, for both preclinical and clinical studies. Indeed since glucan synergizes equally well with IgM antibody, the presence of natural IgM anti-tumor and anti-viral antibodies can be a confounding factor in interpreting in vivo tumor response, whether in preclinical models or in clinical trials, unless the po intake of glucan in mouse chow is standardized. Most importantly, since many carbohydrate tumor vaccines (e.g. GM2-KLH,<sup>35</sup> GD2-KLH, MUC-1,<sup>36</sup> and globo-H-hexasaccharide<sup>37</sup>) induce primarily specific IgM response, glucan may enhance their anti-tumor effects. If this glucan effect can be



generalized to other antibody-mediated host defense mechanisms, its role in infectious disease may also be intriguing. Serotherapy of certain drug resistant bacteria,<sup>38</sup> or viral (e.g. CMV) and fungal (e.g. cryptococcus and candida<sup>39</sup>) infections using antibodies may be enhanced by concurrent intake of  $\beta$ -glucan. One can speculate if the function of pre-existing protective antibodies, e.g. towards tetanus or streptococcus, can be enhanced by oral  $\beta$ -glucan; indeed, if it can enhance the protective effects of common bacterial vaccines. The successful treatment of Alzheimer's disease using antibodies specific for amyloid  $\beta$ -peptide in the mouse model is a provocative finding<sup>44</sup>; it is likely that  $\beta$ -glucan may enhance the antibody effect. When one considers glucan-effect in the context of auto-immune disease, it is also plausible that tissue injury may be increased by oral glucan, leading to exacerbations of such diseases as rheumatoid arthritis. It is possible in those auto-immune diseases in which auto-antibodies cause tissue damage, clinical signs and symptoms may be modulated by oral intake of glucan. In view of these potential beneficial and adverse effects of barley glucan on human diseases, a better understanding of their immune effects seems highly worthwhile.

## REFERENCES

1. DiMaggio JJ, Scheinberg DA, Houghton AN: Monoclonal antibody therapy of cancer. In: Pinedo HM, Chabner BA, Longo DL, (eds.): Cancer Chemotherapy and Biological Response  
5 Modifiers, Annual 11, Elsevier Science Publishers B.V., (Biomedical Division), 1990, pp 177-203
2. Schlom J: Monoclonal Antibodies in cancer therapy: Basic principles. In: DeVita VT, Hellman S, Rosenberg SA, (eds.): Biologic therapy of cancer, 2nd ed. Philadelphia, J.B.Lippincott Co, 1995, pp 507-520  
10
3. Koehler G, Milstein C: Continuous culture of fused  
15 cells secreting antibody of pre-defined specificity. Nature 256:495-496, 1975
4. Moffat R, Pinsky CM, Hammershaimb L, et al: Clinical utility of external immunoscintigraphy with the IMMU-4  
20 technetium-99m Fab' antibody fragment in patients undergoing surgery for carcinoma of the colon and rectum: results of a pivotal, phase III trial. The Immunomedics Study Group. J Clin Oncol 14(8):2295-2305, 1996
- 25 5. Maloney DG, Grillo-Lopez AJ, Bodkin DJ, et al: IDEC-C2B8: Results of a phase I multiple-dose trial in patients with relapsed non-hodgkin's lymphoma. J Clin Oncol 15:3266-3274, 1997
- 30 6. Cobleigh MA, Vogel CL, Tripathy D, et al: Multinational study of the efficacy and safety of humanized anti-HER2 monoclonal antibody in women who have HER2-overexpressing metastatic breast cancer that has progressed after chemotherapy for metastatic disease. J Clin Oncol 17:2639-  
35 2648, 1999
7. Bigner DD, Brown MT, Friedman AH, et al: Iodine-131-labeled antitenascin monoclonal antibody 81C6 treatment of

patients with recurrent malignant gliomas:phase I trial results. Journal Clinical Oncology 16:2202-2212, 1998

8. Jurcic JG, Caron PC, Miller WH, et al: Sequential  
5 targeted therapy for acute promyelocytic leukemia with all-trans retinoic acid and anti-CD33 monoclonal antibody M195. Leuk 9:244-248, 1995
9. Meredith RF, Khazaeli MB, Plott WE: Phase II study of  
10 dual <sup>131</sup>I-labeled monoclonal antibody therapy with interferon in patients with metastatic colorectal cancer. Clin Can Res 2:1811-1818, 1996
10. Yeh SD, Larson SM, Burch L, et al: Radioimmunodetection  
15 of neuroblastoma with iodine-131-3F8: Correlation with biopsy, iodine-131-Metaiodobenzylguanidine (MIBG) and standard diagnostic modalities. J Nucl Med 32:769-776, 1991
11. Cheung NKV, Kushner BH, Cheung IY, et al: Anti-GD2  
20 antibody treatment of minimal residual stage 4 neuroblastoma diagnosed at more than 1 year of age. J Clin Oncol 16:3053-3060, 1998
12. Lanzavecchia A, Abrignani S, Scheidegger D, et al:  
25 Antibodies as antigens. The use of mouse monoclonal antibodies to focus human T cells against selected targets. J Exp Med 167:345-352, 1988
13. Jerne NK: Toward a network theory of the immune system.  
30 Ann Immunol (Paris) 125C:373-389, 1974
14. Janeway CA: Presidential address to the american association of immunologist. J Immunol 161:539-544, 1998
15. Liszewski MK, Farries TC, Lublin DM, et al: Control of  
35 the complement System. Adv Immunol 61:201, 1996

16. Kim YS, Ryu J, Han S, et al: Gram-negative bacteria binding protein, a pattern recognition receptor for lipopolysaccharide and beta-1,3-glucan, which mediates the signaling for the induction of innate immune genes in *Drosophila melanogaster* cells. *J Bio Chem* 2000
17. Bohn JA, BeMiller JN: (1-3)  $\beta$ -D-Glucans as biological response modifiers: a review of structure-functional activity relationships. *Carbohydrate Polymers* 28:3-14, 1995
18. Cheung NK, Saarinen U, Neely J, et al: Monoclonal antibodies to a glycolipid antigen on human neuroblastoma cells. *Cancer Res* 45:2642-2649, 1985
19. Modak S, Gultekin SH, Kramer K, et al: Novel tumor-associated surface antigen: broad distribution among neuroectodermal, mesenchymal and epithelial tumors, with restricted distribution in normal tissues. *Proceedings of ASCO* 17:449a, 1998
20. Nethery A, Raison RL, Easterbrook-Smith SB: Single-step purification of immunoglobulin M on Clq-sepharose. *J Immunol Methods* 126:57-60, 1990
21. Cheung NKV, Walter EI, Smith-Mensah WH, et al: Decay-accelerating factor protects human tumor cells from complement-mediated cytotoxicity in vitro. *J Clin Invest* 81:1122-1128, 1988
22. Houghton AN, Mintzer D, Cordon-Cardo C, et al: Mouse monoclonal antibody detecting GD3 ganglioside: a phase I trial in patients with malignant melanoma. *Proc Natl Acad Sci USA* 82:1242-1246, 1985
23. Munn DH, Cheung NK: Interleukin-2 enhancement of monoclonal antibody-mediated cellular cytotoxicity (ADCC) against human melanoma. *Cancer Res* 47:6600-6605, 1987

24. Zimmerman JW, Lindermuth J, Fish PA, et al: A novel carbohydrate-glycosphingolipid interaction between a beta-(1-3)-glucan immunomodulator, PGG-glucan, and lactosylceramide of human leukocytes. J Biol Chem 273:22014-22020, 1998
25. Matzinger P: Tolerance, danger and the extended family. Annu Rev Immunol 12:991-1045, 1994
26. Fuchs EJ, Matzinger P: Is cancer dangerous to the immune system. Seminars in Immunology 8:271-280, 1996
27. Matzinger P: An innate sense of danger. Seminars in Immunology 10:399-415, 1998
28. Medzhitov R, Janeway CA: An ancient system of host defense. Curr Opin Immunol 10:12-15, 1998
29. Vivanco F, Munoz E, Vidarte L, Pastor C: The covalent interaction of C3 with IgG immune complexes. Mol Immunol 36:843-852, 1999
30. Vetvicka V, Thornton BP, Ross GD: Soluble beta-glucan polysaccharide binding to the lectin site of neutrophil or natural killer cell complement receptor type 3 (CD11b/CD18) generates a primed state of the receptor capable of mediating cytotoxicity of iC3b-opsonized target cells. Journal of Clinical Investigations 98:50-61, 1996
31. Yan J, Vetvicka V, Xia Y, et al: B-glucan a "Specific" biologic response modifier that uses antibodies to target tumors for cytotoxic recognition by leukocyte complement receptor type 3 (CD11b/CD18). J Immunol 163:3045-3052, 1999
32. Xia Y, Vetvicka V, Yan J, et al: The beta-glucan-binding lectin site of mouse CR3 (CD11b/CD18) and its function in generating a primed state of the receptor that

mediates cytotoxic activation in response to iC3b-opsonized target cells. J Immunol 162:2281-2290, 1999

5 33. Thornton BP, Vetvicka V, Ross GD: Function of C3 in a humoral response: iC3b/C3dg bound to an immune complex generated with natural antibody and a primary antigen promotes antigen uptake and the expression of co-stimulatory molecules by all B cells, but only stimulates immunoglobulin synthesis by antigen-specific B cells. Clin Exp Immunol  
10 104:531-537, 1996

34. Jurianz K, Maslak S, Garcia-Schuler H, et al: Neutralization of complement regulatory proteins augments lysis of breast carcinoma cells targeted with rhumAb anti-  
15 HER2. Immunopharmacology 42:209-218, 1999

35. Livingston PO, Wong GYC, Adluri S, et al: Improved survival in stage III melanoma patients with GM2 antibodies: a randomized trial of adjuvant vaccination with GM2 ganglioside. J Clin Oncol 12:1036-1044, 1994  
20

36. Kotera Y, Fontenot JD, Pecher G, et al: Humoral immunity against a tandem repeat epitope of human mucin MUC-1 in sera from breast, pancreatic, and colon cancer  
25 patients. Cancer Res 54:2856-60, 1994

37. Slovin SF, Ragupathi G, Adluri S, et al: Carbohydrate vaccines in cancer: Immunogenicity of a fully synthetic globo H hexasaccharide conjugate in man. Proc Natl Acad Sci  
30 USA 96:5710-5715, 1999

38. Casadevall A: Antibody-based therapies for emerging infectious diseases. Emerging Infectious Diseases 2: 1996

35 39. Han Y, Riesselman MH, Cutler JE: Protection against candidiasis by an immunoglobulin G3 (IgG3) monoclonal

antibody specific for the same mannotriose as an IgM protective antibody. Inf Imm 68:1649-1654, 2000

5 40. Waksal HW: Role of an anti-epidermal growth factor receptor in treating cancer. Cancer and Metastasis Reviews 18:427-436, 1999

10 41. Agus DB, Scher HI, Higgins B, et al: Response of prostate cancer to anti-her-2/neu antibody in androgen-dependent and independent human xenograft models. Advances in Brief 59:4761, 1999

15 42. Scott AM, Geleick D, Rubira M, et al: Construction, production, and characterization of humanized anti-lewis Y monoclonal antibody 3S193 for targeted immunotherapy of solid tumors. Cancer Res 60:3254-3261, 2000

20 43. Clarke K, Lee F, Brechbiel MW, et al: Therapeutic efficacy of anti-Lewis humanized 3S193 radioimmunotherapy in a breast cancer model: enhanced activity when combined with taxol chemotherapy. Clin Can Res 6:3621, 2000

25 44. Bard F, Cannon C, Barbour R, et al: Peripherally administered antibodies against amyloid B-peptide enter the central nervous system and reduce pathology in a mouse model of Alzheimer disease. Nature Medicine 6:916-919, 2000

**Second Series of Experiments**

**Purpose:** In vitro  $\beta$ -glucan can enhance tumor cytotoxicity through iC3b receptors on leukocytes. We test if (1 $\rightarrow$ 3), (1 $\rightarrow$ 4)- $\beta$ -D-glucan ( $\beta$ -glucan) can synergize with anti-GD2 monoclonal antibody (MoAb) 3F8 (mouse IgG3) in therapy of human neuroblastoma xenografts.

**Experimental Design:** Athymic nude mice with established neuroblastoma xenografts were treated with daily intraperitoneal or oral  $\beta$ -glucan, in the presence/absence of intravenous MoAb twice a week, for 22-29 days. Serial tumor volumes and body weights were monitored.

**Results:** 3F8 plus  $\beta$ -glucan produced near-complete tumor regression/disease stabilization, while 3F8 or  $\beta$ -glucan alone did not significantly affect tumor growth. For NMB7 tumors, median survival of 3F8 plus  $\beta$ -glucan group was 5.5 fold that of control groups ( $p < 0.001$ ), and for LAN-1 the survival difference was 2.6 fold. 47% of the mice with NMB7 and 18% with LAN-1 remained progression-free in contrast to  $< 3\%$  of controls. Antitumor effect was seen at 340 ug glucan dose, i.v. or po, and in all human neuroblastoma cell lines tested. No toxicities were noted in mice treated with either  $\beta$ -glucan alone or 3F8 plus  $\beta$ -glucan (4 to 4000 ug per dose). In contrast to anti-GD2 MoAb 3G6 (IgM), 3F8 F(ab')<sub>2</sub> and MoAb 8H9 (IgG1) did not activate complement and had no synergy with  $\beta$ -glucan. Anti-tumor effect of 3F8 plus oral  $\beta$ -glucan persisted after anti-asialo-GM1 antibody treatment, as well as in NK-deficient host.

**Conclusions:** Oral 1,3-1,4- $\beta$ -glucan synergized with anti-tumor IgG and IgM MoAb in vivo. Since  $\beta$ -glucan was well tolerated and inexpensive, its potential value in cancer therapy deserves further investigation.



**INTRODUCTION**

Monoclonal antibodies (MoAb) selective for tumors have therapeutic potential (1). The introduction of hybridoma  
5 technology by Kohler and Milstein in 1975 (2) and advances in molecular biologic techniques have greatly expanded the potential of MoAb in human cancers. Evidence of efficacy in clinical trials is increasingly evident: 17-1A in colon  
cancers (3), anti-CD20 in lymphoma (4, 5), anti-HER2  
10 antibodies in breast cancer (6, 7), and M195 against CD33 in acute leukemia (8) are good examples. Our laboratory has developed the MoAb 3F8 which targets the ganglioside GD2 overexpressed on neuroblastoma. 3F8 has been shown to have high specificity and sensitivity in the radioimmunodetection  
15 of minimal residual disease in patients with NB (9), and a significant clinical impact when used as adjuvant therapy (10).

The immune basis of clinical tumor response to MoAb is at  
20 least two-fold, direct cytotoxicity and induced immunity. Antibody dependent cell-mediated cytotoxicity (ADCC) and complement mediated cytotoxicity (CMC) are responsible for the direct killing of tumor cells. On the other hand, through tumor opsonization (11) or idiotype network (12),  
25 tumor-specific immunity is induced.  $\beta$ -glucans are polymers of glucose extractable from cereals, mushrooms, seaweed and yeasts (13). They are (1 $\rightarrow$ 3)- $\beta$ -D-glucopyranosyl polymers with randomly dispersed single  $\beta$ -D-glucopyranosyl units attached by (1 $\rightarrow$ 6)- linkages, giving a comb-like structure.  
30 The (1 $\rightarrow$ 3)- $\beta$  backbone and the (1 $\rightarrow$ 6)-linked branches were thought to be important for their immune effects. Lentinan (from Lentinus edodes, Basidiomycete family) is a high molecular weight (MW)  $\beta$ -glucan with (1 $\rightarrow$ 6) branches off every three (1 $\rightarrow$ 3)- $\beta$ -D-glucopyranosyl residues and it has been  
35 licensed in Japan for cancer treatment. Schizophyllan (from Schizophyllum commune, Basidiomycete family) and  $\beta$ -glucan

from Baker's yeast (*Saccharomyces cerevisiae*) have similar structures. Laminarin (from seaweed), a small MW  $\beta$ -glucan, has (1 $\rightarrow$ 6)- $\beta$  branches occurring at every ten (1 $\rightarrow$ 3)- $\beta$ -D glucopyranosyl units. On the other hand,  $\beta$ -glucan from  
5 barley, oat or wheat has mixed (1 $\rightarrow$ 3) and (1 $\rightarrow$ 4)- $\beta$ -linkage in the backbone, but no (1 $\rightarrow$ 6)- $\beta$  branches and are generally of high MW. Although barley (1 $\rightarrow$ 3), (1 $\rightarrow$ 4)- $\beta$ -D-glucan has been shown in vitro to bind to CR3 (14), activate ADCC mediated by natural killer cells (15-17), monocytes (18, 19), and  
10 neutrophils (17, 19), as well as stimulating tumor necrosis factor (TNF $\alpha$ ) production by monocytes (20), their in vivo immunomodulatory effects in cancer models have yet to be investigated.

15 We now report our findings that oral (1 $\rightarrow$ 3), (1 $\rightarrow$ 4)- $\beta$ -D-glucan derived from barley or oats can greatly enhance the activity of anti-tumor monoclonal antibodies in xenograft models. Because  $\beta$ -glucan is nontoxic, well tolerated and inexpensive, its role in cancer therapy deserves careful  
20 study.

#### MATERIALS AND METHODS

**Cell lines** Human neuroblastoma cell lines LAN-1 were  
25 provided by Dr. Robert Seeger, Children's Hospital of Los Angeles, Los Angeles, CA, and NMB7 by Dr. Shuen-Kuei Liao (McMaster University, Ontario, Canada). Neuroblastoma cell lines SK-N-JD, SK-N-ER, and SK-N-MM were established from patients with metastatic disease treated at Memorial Sloan-  
30 Kettering Cancer Center (MSKCC), New York, NY. Cell lines were cultured in 10% defined calf serum (Hyclone, Logan, UT) in RPMI with 2 mM L-glutamine, 100 U/ml of penicillin (Sigma, St. Louis, MO), 100 ug/ml of streptomycin (Sigma), 5% CO<sub>2</sub> in a 37°C humidified incubator. Normal human  
35 mononuclear cells were prepared from heparinized bone marrow

samples by centrifugation across a Ficoll density separation gradient.

**Antibodies** Monoclonal antibodies 3F8 (mouse IgG3), 3G6 (mouse IgM), and 8H9 (mouse IgG1) reactive with neuroblastoma have been previously described (21, 22). They were produced as ascites and purified by affinity chromatography: protein A (Pharmacia, Piscataway, NJ) for 3F8 (21), protein G (Pharmacia) for 8H9 (22), and Clq-sepharose (Pierce, Rockford, IL) for 3G6 (21, 23). These antibodies were >90% pure by SDS-PAGE. F(ab')<sub>2</sub> fragments were prepared by pepsin digestion as previously reported (24). TIB114 (N.S.7), a hybridoma secreting an IgG3 control antibody, was obtained from American Type Culture Collection (ATCC). Rabbit anti-asialo-GM1 antibody (Wako Pure Chemical Industries, Osaka, Japan) diluted to 1 mg/ml of protein was administered at 200 ul ip on days 0, 1, 2, 7, 14, 21.

**Glucan** (1→3), (1→4)-β-D-glucan derived from barley and (1→4)-β-D-mannan were purchased from Sigma. Sugar composition and linkage analysis were performed by the Complex Carbohydrate Research Center, University of Georgia, Athens, Georgia, supported in part by the Department of Energy-funded Center for Plant and Microbial Complex Carbohydrates (DF-FG09-93ER-20097). Barley glucan was dissolved by boiling for 10 minutes in normal saline.

**Mice and treatment** Athymic Balb/c mice were purchased from NCI, Frederick, MD, and CB-17 SCID-Beige mice from Taconic (Germantown, NY). Tumor cells were planted (1-5 x10<sup>6</sup> cells) in 100ul of Matrigel (BD BioSciences, Bedford, MA) subcutaneously. Following implantation, tumor sizes (maximum width and lengths) were measured. Tumor size was calculated as product of the 2 perpendicular diameters. Treatment studies started in groups of 4-5 mice per cage

when tumor diameter reached 0.7 to 0.8 cm, usually by 14-21 days of tumor implantation. Mice received antibody treatment intravenously (generally 200 ug per dose by retroorbital injection) twice weekly and  $\beta$ -glucan by intragastric gavage (generally 400 ug per dose) every day for 3 weeks (22-days of  $\beta$ -glucan and 6-doses of antibody). Mice were weighed once a week and tumor size measured twice a week. Mice were sacrificed when tumors reached sizes that interfered with their well-being.

**Assays for soluble cytokines** Sera from mice were obtained 1 h, 4 h, 8 h, 24 h, 48 h, and 72 h after oral  $\beta$ -glucan. They were assayed for soluble cytokine IL-12 (p70) and TNF $\alpha$ , all reagents from Endogen (Woburn, MA). Briefly, 96-well microtiter plates were coated with either monoclonal anti-mouse IL12 at 5 ug/ml or monoclonal anti-TNF $\alpha$  at 0.8 ug/ml overnight at ambient temperature. The mouse IL12 standard ranged from 1000 pg/ml in 1:3 serial dilutions and the TNF $\alpha$  standard ranged from 490 pg/ml in 1:2 serial dilution. Test samples (serum diluted 1:2) were added to the plates and incubated for 2 hours at ambient temperature. The detecting antibody, biotinylated anti-mouse IL12 monoclonal at 1:100 dilution for the IL12 assay, or biotinylated anti-mouse TNF $\alpha$  monoclonal at (1:50) for the TNF $\alpha$  ELISA was added. The plates were incubated at ambient temperature for one hour. After PBS wash, the secondary antibody, which was HRP-conjugated streptavidin at 1:400 for IL12, and 1:200 for TNF $\alpha$ , was added to the plates for a 30 min incubation at ambient temperature. After another wash, tetramethylbenzidine was added as the substrate for the color reaction for 30 min, and absorbance was read at 450 nm using an ELISA plate reader. The limits of detection were 12 pg/ml for the mouse IL12 ELISA, and 10 pg/ml for the mouse TNF $\alpha$  ELISA.

**Immunostaining for tumor vasculature** LAN-1 xenografts were removed 1h, 4h, 8h, 16h, 24h, 48h, 96h and 216 h after treatment. Tumor vasculature was assayed by immunostaining with an anti-blood vessel antibody. Eight mm cryostat  
5 frozen tumor sections were fixed in acetone and washed in PBS. Endogenous peroxidases were blocked in 0.3% H<sub>2</sub>O<sub>2</sub> in PBS. Sections were incubated in 3% bovine serum albumin containing 0.25% gelatin for 60 minutes, after the avid-biotin blocking step. Incubation with the biotinylated rat  
10 anti-murine PECAM IgG2a MoAb, MEC13.3 (1 mg/ml) (BD PharMingen, SanDiego, CA ) was carried out at room temperature for 60 minutes followed by ABC complex ( Vector Laboratories, Burlingame, CA ). Color was developed with DAB peroxidase substrate kit (Vector). A 10% hematoxylin  
15 counterstain for 4 minutes was used.

**Statistical analysis** Average tumor size over time between groups was compared. The null hypothesis was no difference in size over time. To test the hypothesis, the square of  
20 size differences summed over time was used, which in effect compared the trajectories of the average tumor sizes between treatment groups.

$$\text{SS\_DEV} = 3 \sum_{i=1}^k (x_i - y_i)^2$$

where there were k time points and  $x_i$  and  $y_i$  were the average tumor sizes at time i for each treatment group.

30

## RESULTS

**Synergy between barley  $\beta$ -glucan and anti-GD2 antibody 3F8 in eradicating human neuroblastoma.** 3F8 is a murine IgG3  
35 monoclonal antibody that activates mouse and human complement, and mediates effective ADCC against human neuroblastoma cells in vitro.  $\beta$ -glucan when administered

orally at 400 ug per day had no appreciable effect on NMB7 tumor growth as did antibody 3F8 given i.v. alone. However, when  $\beta$ -glucan and 3F8 were used in combination, tumor growth was near totally suppressed. In >47% of mice, tumors remained permanently suppressed following treatment. Similar observations were made with neuroblastoma cell lines derived from different sources: LAN-1 (**Figure 1a**), NMB7 (**Figure 1b**), SK-N-ER (**Figure 1c**), SK-N-MM and SK-N-JD (data not shown).  $\beta$ -glucan was equally effective when administered orally or intraperitoneally. In contrast, for the GD2-negative rhabdomyosarcoma, HTB82, 3F8 plus  $\beta$ -glucan treatment was ineffective (data not shown). In addition, TIB114 (IgG3 control) plus barley  $\beta$ -glucan, or 3F8 plus mannan had no anti-tumor (data not shown). When 3F8 dose was decreased from 200 ug to 40 ug, the anti-tumor effect was lost (data not shown). There was no detectable serum IL-12 or TNF- $\alpha$  release following oral  $\beta$ -glucan administration (data not shown). There was no immunohistochemically detectable effect of  $\beta$ -glucan on tumor vessel formation (data not shown).

**Dose response curve for ip  $\beta$ -glucan.** When the dose of intraperitoneal  $\beta$ -glucan was decreased by 10-fold from 4000 ug, it was clear that 4 ug was no longer effective in synergizing with MoAb 3F8 in suppressing NMB7 growth. Interestingly, both ip and oral 1,3-1,4- $\beta$ -glucan (at 400 ug per day) were effective (**Figure 2**).

**Oral  $\beta$ -glucan versus ip  $\beta$ -glucan.** When oral  $\beta$ -glucan was studied in NMB7 tumors (**Figure 3**), similar dose response was found. While an oral dose of 400 ug was curative for some tumors, breakthroughs were seen for lower dose levels, with those receiving 4 ug escaping sooner than those receiving 40 ug. Using the LAN-1 tumor model, neither 4 nor 40 ug  $\beta$ -glucan were effective (data not shown). There was no significant

body weight change in any of the treatment groups (after accounting for tumor weight), irrespective of  $\beta$ -glucan dose or co-administration with 3F8. At necropsy on day 22, there were no appreciable differences in the peripheral blood counts, cholesterol and blood chemistry between mice receiving different  $\beta$ -glucan doses. There was also no difference in the histologic appearances of organs in mice treated with  $\beta$ -glucan at any of the dose levels, when compared to control mice that received saline.

By the oral route, daily  $\beta$ -glucan schedule was necessary. A 5 day/week po  $\beta$ -glucan regimen was comparable to the daily regimen. In contrast, a once a week or twice a week schedule of  $\beta$ -glucan had no anti-tumor effect (data not shown).

**Role of NK cells in  $\beta$ -glucan effect.** Removal of NK cells by anti-Asialo GM1 antiserum eliminated a substantial amount, although not completely the anti-tumor activity of  $\beta$ -glucan (Figure 4). Moreover, in SCID-beige mice which lack NK cells,  $\beta$ -glucan was still effective in synergizing with 3F8 (data not shown), suggesting that at least part of the anti-tumor activity was mediated by NK-independent cytotoxicity.

**IgG3-F(ab')<sub>2</sub> or IgG1 antibodies did not have anti-tumor activity.** The role of Fc in mediating the anti-tumor effect of  $\beta$ -glucan was apparent when Fc was removed by pepsin or when IgG1 isotype MoAb 8H9 was used (data not shown). Neither was able to activate complement or mediate efficient ADCC, and neither has significant anti-tumor effect when administered with 400 ug of oral  $\beta$ -glucan.

**Synergy of  $\beta$ -Glucan with 3F8 in prolonging survival.** Nude mice (n=22) with established neuroblastoma NMB7 xenografts (0.7-0.8 cm diameter tumor at the beginning of treatment)

were treated with 3F8 (200 ug twice a week iv) and 400 ug of  $\beta$ -glucan po daily for a total of 3 weeks. Control mice received either saline alone (n=10), 3F8 alone (n=8), or  $\beta$ -glucan (n=16) alone. Median survival was 30 days in control groups and 166 days in the treatment (3F8 plus  $\beta$ -glucan, n=22) group ( $p < 0.001$ ). Long-term survival was estimated at 47% in the treatment group and 3% in the control group (saline alone, 3F8 alone, or  $\beta$ -glucan alone) (**Figure 5**). Similar experiments were carried out in nude mice bearing established LAN-1 xenografts (also 0.7-0.8 cm diameter tumor at the beginning of treatment). Among control mice treated with either saline alone (n=31), 3F8 alone (n=16), or  $\beta$ -glucan (n=8) alone, tumor growth was rapid. Median survival was 21 days in control groups (n=55) and 54 days in the treatment (3F8 plus  $\beta$ -glucan) group (n=82) ( $p < 0.001$ , **Figure 6**). Long-term survival was estimated at 18% in the treatment group and 0% in the controls.

## DISCUSSION

Using the human xenograft models, we have made the following observations.  $\beta$ -Glucan derived from barley can synergize with monoclonal antibodies to suppress and/or eradicate tumors, while  $\beta$ -glucan or antibody alone has little anti-tumor effect. Anti-tumor response requires antibodies that activate complement, and both mouse IgM and mouse IgG3 were effective. Oral administration of  $\beta$ -glucan is at least equally (if not more) effective than the intraperitoneal route. It is a dose-dependent phenomenon, where  $\geq 400$  ug per dose is required for maximal effect. Natural killer cells are not essential for this  $\beta$ -glucan phenomenon, although they contribute to the anti-tumor effect. Normal T-cells and B-cells are not required for the anti-tumor effect since immune-deficient mouse strains demonstrate the  $\beta$ -glucan effect, in both athymic and SCID-beige mice. Most importantly, oral  $\beta$ -glucan is well-tolerated by all the mice



tested so far, with no noticeable change in body weight, blood counts or organ histologies, even at doses as high as 4 mg per dose per day.

5 Our findings differ significantly from previous observations and predictions on the use of  $\beta$ -glucans in cancer treatment. In the past it was thought that the (1 $\rightarrow$ 3), (1 $\rightarrow$ 6)- $\beta$  linkage was absolutely required for the  $\beta$ -glucan anti-tumor effect (13). This structure contains (1 $\rightarrow$ 3)- $\beta$ -D-glucopyranosyl  
10 units along which are randomly dispersed single  $\beta$ -D-glucopyranosyl units attached by (1 $\rightarrow$ 6)- linkages, giving a comb-like structure (e.g. Lentinan, Schizophyllan, Laminarin, and glucan from Baker's yeast). In these models, it was believed that T-cells were activated and  
15 indeed required for the anti-tumor effect. In addition, it was believed that small molecular weight  $\beta$ -glucan should be more effective than high molecular weight  $\beta$ -glucan and that the most effective administration should be intravenous or intraperitoneal routes. Indeed, Betafectin was derived from  
20 a genetically engineered *Saccharomyces cerevisiae* which makes (1 $\rightarrow$ 3), (1 $\rightarrow$ 4)- $\beta$ -D-glucans with weaker interchain associations (25). It was manufactured for i.v. injection to improve macrophage function in the hope of reducing infectious complications and improving wound healing.  
25 Barley  $\beta$ -glucan is a linear polymer with (1 $\rightarrow$ 3)- $\beta$  and (1 $\rightarrow$ 4)- $\beta$  linkages; however, it is not a comb like structure. We did not find any anti-tumor effect of barley  $\beta$ -glucan when given alone. However, when used in combinations with monoclonal antibodies, the synergistic effect was remarkable. Although  
30  $\beta$ -glucan activates granulocyte mediated ADCC in vitro (data not shown), the effects of  $\beta$ -glucan may be indirect. It is not clear if the absorption of  $\beta$ -glucan is necessary for its anti-tumor effect. The exact mechanism of how  $\beta$ -glucan enhances the anti-tumor effect of monoclonal antibodies in  
35 vivo is unknown.

Monoclonal antibodies, either through Fc interaction or through CR3 interaction with iC3b, target cytotoxicity to tumor cells, a process greatly enhanced by  $\beta$ -glucan activation of effector cells.  $\beta$ -glucan activates leukocytes by binding to CR3 or to  $\beta$ -glucan receptors (26). After antibodies deposit complement components on pathogens or cancer cells, C3b is rapidly proteolyzed into iC3b fragment by serum factor I. These iC3b fragments then opsonize the pathogens or tumor cells for the iC3b-receptors (CR3, CD11b/CD18) on phagocytic cells and NK cells, stimulating phagocytosis and/or cytotoxic degranulation. Thus, antibody and complement link innate and adaptive immunity by targeting antigens to different cells of the immune system, e.g. via CR3 and Fc for phagocytic cells, CR2 for B cells, and CR1, CR2, or CR3 for follicular dendritic cells (27). For neutrophils, CR3-dependent phagocytosis requires ligation of two distinct binding sites, one for iC3b and a second site for  $\beta$ -glucan. Without  $\beta$ -glucan, iC3b-opsonized target cells are resistant to killing (17). Microbes possess polysaccharides that can activate the lectin domain on CR3, leading to phagocytosis and cytotoxic degranulation. In contrast, human cells (including tumors) lack these CR3-binding polysaccharides, thus the inability of CR3 to mediate phagocytosis or extracellular cytotoxicity of tumor cells opsonized with iC3b. The lectin site of CR3 can also influence transmembrane signaling of endogenous neutrophil membrane GPI-anchored glycoproteins (e.g. CD14, CD16, and CD59). In a mouse mammary tumor model, where there are naturally occurring IgM and IgG antibodies, injection of yeast soluble  $\beta$ -glucan could suppress tumor growth, an effect lost in C3-deficient or CD11b (CR3)-deficient mice (28, 29). Since iC3b bound to a primary protein antigen can also enhance recognition and specific antibody synthesis by antigen-specific B cells (30), the

presence of  $\beta$ -glucan plus complement activation may enhance B-cell response to pathogens or tumor cells.

Another mechanism of action of  $\beta$ -glucan may relate to innate  
5 receptors for  $\beta$ -glucan, in a hard-wired information network  
on phagocytes and lymphoid cells; receptors that normally  
recognize death signals and microbial molecular patterns  
(31). These innate receptors are biosensors for invading  
10 pathogens widely distributed in vertebrate and invertebrate  
animals (32), a nonclonal host defense pathway with  
structural and functional homologies in phylogenetic  
lineages that diverged over a billion years ago. Following  
 $\beta$ -glucan activation of leukocytes, killing is immediate,  
15 nonclonal, and obligatory, a process often referred to as  
innate immunity. The consequence of this innate effector  
arm is the activation of costimulatory molecules and  
induction of cytokines and chemokines that will enhance  
adaptive immunity to the tumor cells. (33, 34). Thus,  
20 antibodies, complement, phagocytes, and "danger" receptors  
form core elements of innate immunity while  
antigen-presenting cells, T and B lymphocytes constitute  
essential players in acquired immunity.

Despite the availability of tumor-selective monoclonal  
25 antibodies and the ample supply of phagocytes/natural  
killers, shrinkage of established tumors following antibody  
treatment alone, and the acquisition of specific immunity,  
are not common in both preclinical models and cancer  
patients. The absence of a danger signal and the diminution  
30 of complement action by complement resistance proteins on  
tumor cells may explain the inefficiency of antibody  
mediated clinical responses (35). Lipopolysaccharide and  
 $\beta$ -glucan, being cell wall components of bacteria and fungus,  
respectively, are potent danger signals to the immune  
35 systems in all life-forms, from *Drosophila* to man (36).

While LPS is too toxic for human use,  $\beta$ -glucan is relatively benign. If this synergistic effect of  $\beta$ -glucan on antibodies is active in humans, our findings may have broad clinical implications. First the efficacy of monoclonal  
5 antibodies in cancer (e.g. Herceptin, Rituximab, Dacluzimab, anti-GD2 and anti-EGF-R MoAb) can be potentially enhanced (37). Nevertheless, even though toxicity from  $\beta$ -glucan is expected to be minimal, the enhanced efficacy of MoAb may also increase MoAb-mediated toxicity. For example, the side  
10 effects of Herceptin on cardiac function, or anti-GD2 MoAb on neuropathic pain may be increased. Second, since the amount and quality of barley and oat glucan in daily food intake can vary, future interpretations of efficacy trials using MoAb may need to take this into account, for both  
15 preclinical and clinical studies. Indeed since glucan synergizes equally well with IgM antibody, the presence of natural IgM anti-tumor and anti-viral antibodies can be a confounding factor in the interpretation of in vivo tumor response, whether in preclinical models or in clinical  
20 trials, unless the oral intake of glucan in mouse chow is standardized. Most importantly, since many carbohydrate tumor vaccines (e.g. GM2-KLH (38), GD2-KLH, and globo-H-hexasaccharide (39)) primarily induce specific IgM response, glucan may enhance their anti-tumor effects. In view of  
25 these potential beneficial effects of barley glucan on cancer therapy, a better understanding of their immune effects seems highly worthwhile.

**REFERENCES**

1. Schlom, J.: Monoclonal Antibodies in cancer therapy: Basic principles. In: V. T., DeVita, S., Hellman, S. A., Rosenberg (eds.), Biologic therapy of cancer, 2nd ed., pp. 507-520. Philadelphia: J.B.Lippincott Co, 1995.
2. Koehler, G., Milstein, C. Continuous culture of fused cells secreting antibody of pre-defined specificity. Nature, 256:495-496, 1975.
3. Riethmuller, G., Holz, E., Schlimok, G., Schmiegeler, W., Raab, R., Hoffken, K., Gruber, R., Funke, I., Pichlamaier, H., Hirche, H., Buggisch, P., Witte, J., Pichlmayr, R. Monoclonal Antibody Therapy for Resected Dukes' C Colorectal Cancer: Seven-Year Outcome of a Multicenter Randomized Trial. J Clin Oncol, 16:1788-1794, 1998.
4. Maloney, D. G., Grillo-Lopez, A. J., Bodkin, D. J., White, C. A., Liles, T.-M., Royston, I., Varns, C., Rosenberg, J., Levy, R. IDEC-C2B8: Results of a phase I multiple-dose trial in patients with relapsed non-hodgkin's lymphoma. J Clin Oncol, 15:3266-3274, 1997.
5. McLaughlin, P., Grillo-Lopez, A. J., Kink, B. K., Levy, R., Czuczman, M. S., Williams, M. E., Heyman, M. R., Bence-Bruckler, I., White, C. A., Cabanillas, F., Jain, V., Ho, A. D., Lister, J., Wey, K., Shen, D., Dallaire, B. K. Rituximab chimeric anti-CD20 monoclonal antibody therapy for relapsed indolent lymphoma: half of patients respond to four-dose treatment program. J Clin Oncol, 16:2825-2833, 1998.
6. Cobleigh, M. A., Vogel, C. L., Tripathy, D., Robert, N. J., Scholl, S., Fehrenbacher, L., Wolter, J. M., Paton, V., Shak, S., Lieberman, G., Slamon, D. J. Multinational study of the efficacy and safety of humanized anti-HER2 monoclonal antibody in women who have HER2-overexpressing metastatic

breast cancer that has progressed after chemotherapy for metastatic disease. J Clin Oncol, 17:2639-2648, 1999.

5 7. Slamon, D. J., Leylan-Jones, B., Shak, S., Fuchs, H., Paton, V., Bajamonde, A., Fleming, T., Eiermann, W., Wolter, J., Pegram, M., Baselga, J., Norton, L. Use of chemotherapy plus a monoclonal antibody against HER2 for metastatic breast cancer that overexpress HER2. N Engl J Med, 344:783, 2001.

10 8. Jurcic, J. G., DeBflasio, T., Dumont, L., Yah, T. J., Scheinberg, D. A. Molecular remission induction with retinoic acid and anti-CD33 monoclonal antibody HuM195 in acute promyelocytic leukemia. Clin Cancer Res, 6:372-380, 15 2000.

9. Yeh, S. D., Larson, S. M., Burch, L., Kushner, B. H., LaQuaglia, M., Finn, R., Cheung, N. K. V. Radioimmunodetection of neuroblastoma with iodine-131-3F8: 20 Correlation with biopsy, iodine-131-Metaiodobenzylguanidine (MIBG) and standard diagnostic modalities. J. Nucl. Med., 32:769-776, 1991.

10. Cheung, N. K. V., Kushner, B. H., Cheung, I. Y., 25 Canete, A., Gerald, W., Liu, C., Finn, R., Yeh, S. J., Larson, S. M. Anti-GD2 antibody treatment of minimal residual stage 4 neuroblastoma diagnosed at more than 1 year of age. J Clin Oncol, 16:3053-3060, 1998.

30 11. Lanzavecchia, A., Abrignani, S., Scheidegger, D., Obrist, R., Dorken, B., Moldenhauer, G. Antibodies as antigens. The use of mouse monoclonal antibodies to focus human T cells against selected targets. J. Exp. Med., 167:345-352, 1988.

35 12. Jerne, N. K. Toward a network theory of the immune system. Ann. Immunol. (Paris), 125C:373-389, 1974.

13. Bohn, J. A., BeMiller, J. N. (1-3)-B-D-Glucans as biological response modifiers: a review of structure-functional activity relationships. *Carbohydr Polymers*, 28:3-14, 1995.
14. Xia, Y, Vetvicka, V., Yan, J., Hanikyrova, M., Mayadas, T., Ross, G. D. The beta-glucan-binding lectin site of mouse CR3 (CD11b/CD18) and its function in generating a primed state of the receptor that mediates cytotoxic activation in response to iC3b-opsonized target cells. *J Immunol*, 162:2281-2290, 1999.
15. Di Renzo, L., Yefenof, E., Klein, E. The function of human NK cells is enhanced by beta-glucan, a ligand of CR3 (CD11b/CD18). *Eur J Clin Nutr*, 21:1755-1758, 1991.
16. Vetvicka, V., Thornton, B. P., Wieman, T. J., Ross, G. D. Targeting of natural killer cells to mammary carcinoma via naturally occurring tumor cell-bound iC3b and beta-glucan-primed CR3 (CD11b/CD18). *J Immunol*, 159:599-605, 1997.
17. Vetvicka, V., Thornton, B. P., Ross, G. D. Soluble beta-glucan polysaccharide binding to the lectin site of neutrophil or natural killer cell complement receptor type 3 (CD11b/CD18) generates a primed state of the receptor capable of mediating cytotoxicity of iC3b-opsonized target cells. *Journal of Clinical Investigations*, 98:50-61, 1996.
18. Czop, J. K., Austen, K. F. Properties of glycans that activate the human alternative complement pathway and interact with the human monocyte beta-glucan receptor. *J Immunol*, 135:3388-3393, 1985.
19. Thornton, B. P., Vetvicka, V., Pitman, M., Goldman, R. C., Ross, G. D. Analysis of the sugar specificity and

molecular location of the B-glucan-binding lectin site of complement receptor type 3 (CD11b/CD18). *J Immunol*, 156:1235-1246, 1996.

- 5 20. Roubroeks, J. P., Skjak-Braek, G., Ryan, L., Christensen, B. E. Molecular weight dependency on the production of the TNF stimulated by fractions of rye (1-->3), (1-->4)-beta-D-glucan. *Scand J Immunol*, 52:584-587, 2000.
- 10 21. Cheung, N. K., Saarinen, U., Neely, J., Landmeier, B., Donovan, D., Coccia, P. Monoclonal antibodies to a glycolipid antigen on human neuroblastoma cells. *Cancer Res*, 45:2642-2649, 1985.
- 15 22. Modak, S., Kramer, K., Humayun, G., Guo, H. F., Cheung, N. K. V. Monoclonal antibody 8H9 targets a novel cell surface antigen expressed by a wide spectrum of human solid tumors. *Cancer Res*, 61:4048-4054, 2001.
- 20 23. Nethery, A., Raison, R. L., Easterbrook-Smith, S. B. Single-step purification of immunoglobulin M on Clq-sepharose. *J Immunol Methods*, 126:57-60, 1990.
- 25 24. Cheung, N. K. V., Walter, E. I., Smith-Mensah, W. H., Ratnoff, W. D., Tykocinski, M. L., Medof, M. E. Decay-accelerating factor protects human tumor cells from complement-mediated cytotoxicity in vitro. *J Clin Invest*, 81:1122-1128, 1988.
- 30 25. Zimmerman, J. W., Lindermuth, J., Fish, P. A., Palace, G. P., Stevenson, T. T., DeMong, D. E. A novel carbohydrate-glycosphingolipid interaction between a beta-(1-3)- glucan immunomodulator, PGG-glucan, and lactosylceramide of human leukocytes. *J. Biol. Chem.*, 273:22014-22020, 1998.
- 35



26. Mueller, A., Raptis, J., Rice, P. J., Kalbfleisch, H., Stout, R. D., Ensley, H. E., Browder, W., Williams, D. L. The influence of glucan polymer structure and solution conformation on binding to (1->3)-B-D-glucan receptors in a human monocyte-like cell line. *Glycobiology.*, 10(4):339-346, 2000.
27. Vivanco, F., Munoz, E., Vidarte, L., Pastor, C. The covalent interaction of C3 with IgG immune complexes. *Mol Immunol*, 36:843-852, 1999.
28. Yan, J., Vetvicka, V., Xia, Y., Coxon, A., Carroll, M. C., Mayadas, T. N., Ross, G. D. B-glucan a "Specific" biologic response modifier that uses antibodies to target tumors for cytotoxic recognition by leukocyte complement receptor type 3 (CD11b/CD18). *J Immunol*, 163:3045-3052, 1999.
29. Xia, Y., Vetvicka, V., Yan, J., Hanikyrova, M., Mayadas, T., Ross, G. D. The beta-glucan-binding lectin site of mouse CR3 (CD11b/CD18) and its function in generating a primed state of the receptor that mediates cytotoxic activation in response to iC3b-opsonized target cells. *J Immunol*, 162:2281-2290, 1999.
30. Thornton, B. P., Vetvicka, V., Ross, G. D. Function of C3 in a humoral response: iC3b/C3dg bound to an immune complex generated with natural antibody and a primary antigen promotes antigen uptake and the expression of co-stimulatory molecules by all B cells, but only stimulates immunoglobulin synthesis by antigen-specific B cells. *Clin Exp Immunol*, 104:531-537, 1996.
31. Matzinger, P. Tolerance, danger and the extended family. *Annu Rev Immunol*, 12:991-1045, 1994.

32. Medzhitov, R., Janeway, C. A. An ancient system of host defense. *Curr Opin Immunol*, 10:12-15, 1998.
33. Fuchs, E. J., Matzinger, P. Is cancer dangerous to the  
5 immune system. *Seminars in Immunology*, 8:271-280, 1996.
34. Matzinger, P. An innate sense of danger. *Seminars in Immunology*, 10:399-415, 1998.
- 10 35. Liszewski, M. K., Farries, T. C., Lublin, D. M., Rooney, I. A., Atkinson, J. P. Control of the complement System. *Adv Immunol*, 61:201, 1996.
- 15 36. Kim, Y. S., Ryu, J., Han, S., Choi, K., Nam, K., Jang, I., Lemaitre, B., Brey, P. T., Lee, W. Gram-negative bacteria binding protein, a pattern recognition receptor for lipopolysaccharide and beta-1,3-glucan, which mediates the signaling for the induction of innate immune genes in *drosophila melanogaster* cells. *J Biol Chem*, 2000.
- 20 37. Jurianz, K., Maslak, S., Garcia-Schuler, H., Fishelson, Z., Kirschfink, M. Neutralization of complement regulatory proteins augments lysis of breast carcinoma cells targeted with rhumAb anti-HER2. *Immunopharmacology*, 42:209-218, 1999.
- 25 38. Livingston, P. O., Wong, G. Y. C., Adluri, S., Tao, Y., Padavan, M., Parente, R., Hanlon, C., Calves, M. J., Helling, F., Ritter, G., Oettgen, H. F., Old, L. J. Improved survival in stage III melanoma patients with GM2 antibodies: a randomized trial of adjuvant vaccination with GM2 ganglioside. *J Clin Oncol*, 12:1036-1044, 1994.
- 30 39. Slovin, S. F., Ragupathi, G., Adluri, S., Ungers, G., Terry, K., Kim, S., Spassova, M., Bornmann, W. G., et al. Carbohydrate vaccines in cancer: Immunogenicity of a fully  
35 synthetic globo H hexasaccharide conjugate in man. *Proc. Natl. Acad. Sci. USA*, 96:5710-5715, 1999.

### Third Series of Experiments

**Background:**  $\beta$ -glucan primes leukocyte CR3 for enhanced cytotoxicity and synergizes with anti-tumor monoclonal antibodies (MoAb). We studied (1 $\rightarrow$ 3)- $\beta$ -D-glucans in xenograft tumor models, and examined the relationship of its anti-tumor effect and physico-chemical properties.

**Methods:** Established subcutaneous human xenografts were treated with  $\beta$ -glucan daily and MoAb twice weekly by intragastric injection for 29 days. Control mice received either MoAb alone or  $\beta$ -glucan alone. Tumor sizes were monitored over time.  $\beta$ -glucans were studied by carbohydrate linkage analysis, and high performance size-exclusion chromatography with multiple angle laser scattering detection.

**Results:** Orally administered  $\beta$ -D-glucan greatly enhanced the anti-tumor effects of MoAb against established tumors in mice. This effect correlated with the molecular size of the (1 $\rightarrow$ 3), (1 $\rightarrow$ 4)- $\beta$ -D-glucan. (1 $\rightarrow$ 3), (1 $\rightarrow$ 6)- $\beta$ -D-glucans also synergized with MoAb, although the effect was generally less. We observed this  $\beta$ -glucan effect irrespective of antigen (GD2, GD3, CD20, epidermal growth factor-receptor, HER-2), human tumor type (neuroblastoma, melanoma, lymphoma, epidermoid carcinoma and breast carcinoma) or tumor sites (subcutaneous versus systemic).

**Conclusion:** Given the favorable efficacy and toxicity profile of oral  $\beta$ -D-glucan, its role in cancer treatment as an enhancer of the effect of MoAb therapy deserves careful study.

### **Introduction**

Evidence of efficacy of monoclonal antibodies (MoAb) against human cancer in clinical trials is increasingly evident.

However, induced or administered antibodies to human tumors have not realized their fullest therapeutic potential, even when they can activate complement-dependent cytotoxicity (CDC) and antibody-dependent cell-mediated cytotoxicity (ADCC). The deposition of C3b and iC3b on tumor cells fails to stimulate phagocytosis or extracellular cytotoxicity by C3-receptor-bearing neutrophils, macrophages, and NK cells, even though these same effector cells can efficiently kill C3b and iC3b opsonized microorganisms. The receptor for iC3b, CR3 (also called CD11b/CD18, Mac-1, or  $\alpha M\beta 2$ -integrin), is found in monocytes/macrophages, NK cells, and immune cytotoxic T-lymphocytes. CR3 activation requires the engagement of two sites on its  $\alpha$ -subunit (CD11b): the iC3b-binding site within the I-domain at the N-terminus and a lectin site at the C-terminus (1,2).  $\beta$ -glucans are specific for the lectin site. When coated with iC3b, yeast cells (with their  $\beta$ -glucan-containing cell wall), engage both iC3b and lectin binding sites on leukocytes, triggering phagocytosis and respiratory burst (2,3). In contrast, tumor cells coated with iC3b cannot activate leukocytes because they lack the CR3-binding  $\beta$ -glucan (4-7). Soluble forms of  $\beta$ -glucans can bind to the lectin site (8,9) and prime both phagocytic and NK cells to kill iC3b coated tumor targets (4,9,10). In murine mammary tumor models in which iC3b was found, intravenous yeast  $\beta$ -glucan reduced tumor size by 70-95% (11). The loss of tumor response in the absence of complement-fixing IgM anti-tumor antibodies (SCID mice), or of C3 (C3 knockout mice), or of leukocyte CR3 (CR3 knockout mice) highlighted the critical components of this iC3b strategy (11).

Although (1 $\rightarrow$ 3)- $\beta$ -D-glucans can be purified from yeast, seaweed and mushrooms, an inexpensive, convenient and safe source of pure (1 $\rightarrow$ 3),(1 $\rightarrow$ 4)- $\beta$ -D-glucan is available from barley. Barley  $\beta$ -glucan has been shown to bind to CR3 in

vitro (9), to activate ADCC mediated by NK cells (4,10,12), monocyte (8,13), and neutrophils (8,14), as well as to stimulate TNF production by monocytes(15). However, its in vivo immunomodulatory effects, especially when administered by the oral route, have not been tested. In this study we report an unusually strong synergism between anti-tumor antibodies and intragastric injection of  $\beta$ -glucan against a broad spectrum of human tumor xenografts. We also undertake a preliminary investigation of molecular size requirements for this anti-tumor synergy with MoAb.

### Methods

**Cell lines** The cell lines Daudi, RPMI 6666, HS455, SKMel-28 and A431 were from American Type Culture Collection (ATCC), Rockville, MD. LAN-1 was provided by Dr. R. Seeger, Children's Hospital of Los Angeles, Los Angeles, CA; NMB7 by Dr. S.K. Liao ( McMaster University, Ontario, Canada); human breast carcinoma cell line BT474 was kindly provided by Dr. David Solit of Memorial Sloan-Kettering Cancer Center (MSKCC), New York, NY; SKNJD and SKNER were established at MSKCC. BT474 was cultured in Dulbecco's modified Eagle with Nutrient Mixture F12 (DMEM/F-12) (Life Technologies Inc. GIBCO-BRL, Rockville, MD) in a 1:1 mixture fortified with 10% newborn calf serum (Hyclone, Logan, UT), MEM non-essential amino acids (Gibco-BRL, Grand Island, NY), 100 U/ml of penicillin (Sigma, St. Louis, Mo), and 100 ug/ml of streptomycin (Sigma). All other cell lines were cultured in RPMI 1640 (Life Technologies Inc.) containing 10% defined calf serum (Hyclone) and 100U/ml of penicillin, 100 ug/ml of streptomycin and 2 mM L-glutamine (Sigma).

**Antibodies** MoAb 3F8 (mouse IgG3) and 3G6 (mouse IgM) reactive against GD2 ganglioside expressed on neuroectodermal tumors, and MoAb 8H9 (mouse IgG1) reactive with a glycoprotein expressed on these same tumors have been previously described (16,17). They were purified to >90%

purity by affinity chromatography: protein A (Pharmacia, Piscataway, NJ) for 3F8, and protein G (Pharmacia) for 8H9. Anti-GD3 antibody (R24) (18) was provided by Dr. P. Chapman of MSKCC. Hybridomas producing the anti-epidermal growth factor receptor (EGF-R) antibodies 528 (IgG2a) and 455 (IgG1) were obtained from ATCC (19). Rituximab (anti-CD20) and Herceptin (anti-HER2) were purchased from Genentech, San Francisco, CA.

**$\beta$ -Glucan** Barley, oat and lichenan  $\beta$ -D-glucans were purchased from Sigma and Megazyme International Ireland Ltd., Wicklow, Ireland. Wheat  $\beta$ -glucan was kindly provided by Dr. P. Wood of Agriculture and Agri-Food Canada, West Guelph, Ontario. Betatrim (Quaker Oatrim, 5%  $\beta$ -glucan from oat) was provided by Rhodia Food, Cranbury, NJ. Laminarin was purchased from Sigma and from TCI America, Portland, OR. Lentinan ( $\beta$ -glucan extracted from the mushroom *Lentinus edodes*) was provided by the Drug Synthesis and Chemistry Branch, Developmental Therapeutics Program, Division of Cancer Treatment and Diagnosis, National Cancer Institute, Bethesda, MD.  $\beta$ -glucan was dissolved by boiling for 10 minutes in normal saline; Lentinan was dissolved first in DMSO before diluting in water. Digestion with lichenase (endo-1,3:1,4- $\beta$ -D-glucanase) from *B. subtilis* (Megazyme), was carried out in sodium phosphate buffer (20 mM, pH 6.5) at 40°C for 10 minutes. Sugar composition and linkage analysis by gas chromatography-mass spectrometry following methylation was performed by the Complex Carbohydrate Research Center, University of Georgia, Athens, GA, supported in part by the Department of Energy-funded Center for Plant and Microbial Complex Carbohydrates (DF-FG09-93ER-20097) (20). The average ratio of (1 $\rightarrow$ 3) to (1 $\rightarrow$ 4)- $\beta$ -linkage in (1 $\rightarrow$ 3), (1 $\rightarrow$ 4)- $\beta$ -D-glucans derived from barley, oat and wheat was around 3:7. For molecular size and shape estimations,  $\beta$ -glucan was analyzed by size-exclusion chromatography plus multiple-angle laser light scattering

(MALLS) as previously described (21,22). Besides measuring molecular size by MALLS, the slope derived from root mean square radius versus molar-mass plots gave an estimate of the molecular shape: a slope of 0.33 being the shape of a sphere, 0.5 being random coils and 1.0 being rigid rods. High MW  $\beta$ -glucans were found to be random coils in contrast to low MW species which were more sphere-like.

**Mice and treatment** Athymic nu/nu mice were purchased from the National Cancer Institute-Frederick Cancer Center (Bethesda, Maryland) and ICR/SCID from Taconic (White Plains, NY) and maintained in ventilated cages. Experiments were carried out under Institutional Animal Care and Use Committee (IACUC) approved protocols, and institutional guidelines for the proper and humane use of animals in research were followed. Tumor cells were planted (1-5  $\times 10^6$  cells) in 100  $\mu$ l of Matrigel (Sigma) subcutaneously. Tumor dimensions were measured two to three times a week with vernier calipers, and tumor size was calculated as the product of the two largest perpendicular diameters. For breast carcinoma xenograft studies, 6-8 week old female nude mice (NCI) were initially implanted with 0.72 mg 90-day release 17 $\beta$ -estradiol pellet (Innovative Research of America, Sarasota, FL) subcutaneously into the right flank. Twenty-four hours later,  $10^7$  BT474 cells were implanted subcutaneously into the left flank. All treatment studies started in groups of 4-5 mice when tumor diameters reached 0.7 to 0.8 cm. Mice received antibody treatment (40-200 ug per day) i.v. (by retroorbital injection) twice weekly and oral  $\beta$ -glucan (400 ug per day) by intragastric injection every day for a total 4 weeks. Mice were weighed once a week and sacrificed according to IACUC guidelines. In the SCID mouse systemic human lymphoma (Daudi) model, 5 million cells were administered i.v., and treatment started 10 days later.

**Statistical analysis** Because measurement times varied between experiments, and mice in control groups frequently were sacrificed [as required by IACUC for rapidly enlarging tumors] before the end of each experiment, tumor growth was calculated by fitting a regression slope for each individual mouse to log transformed values of tumor size. Slopes were compared between groups using linear regression with monoclonal antibody treatment,  $\beta$ -glucan treatment and combination treatment as covariates. In the study of melanoma tumor growth,  $\beta$ -glucan was given at two different doses. Thus, dose was added as a covariate for analysis. In the study of epidermoid tumor growth, monoclonal antibody was given at three different doses and antibody dose was added as a covariate. Trends for slope by molecular weight were tested by linear regression of slope scores. Survival analysis was conducted by Cox regression using the indicator variables: monoclonal antibody treatment,  $\beta$ -glucan treatment, and combination treatment; in the survival study of lymphoma, Rituximab was given at two different doses and so dose of antibody was added as a covariate for analysis. All analysis were conducted using STATA (Stata Corporation, College Station, TX).

## Results

Synergy between MoAb and barley  $\beta$ -glucan. We chose the mouse model because of its relative inefficiency in CDC and ADCC (23), and MoAb alone were typically ineffective against established tumors. Oral administration of  $\beta$ -glucan (average MW 210 kD) from barley alone at 400 ug qd x 29 days or antibody 3F8 i.v. alone had no appreciable effect on neuroblastoma LAN-1 tumor growth. The tumor growth rates for the  $\beta$ -glucan alone, 3F8 alone, and saline controls were virtually identical. In contrast, when we combined oral  $\beta$ -glucan with i.v. 3F8, significantly less tumor growth was observed in the 3F8 antibody alone group, 0.7% vs 5.4%



increase in tumor size per day, respectively. In the regression model, only combination treatment significantly reduced tumor growth (4.9% per day, 95% CI 2.4%, 7.4%,  $p=0.001$ ). Nude mice ( $n=22$ ) with established NMB7 xenografts were treated with 3F8 (200 ug twice a week iv) and 400 ug of  $\beta$ -glucan po daily for a total of 4 weeks. Control mice received either saline alone, 3F8 alone, or  $\beta$ -glucan alone. Median survival was 30 days in control groups and 166 days in the treatment (3F8 plus  $\beta$ -glucan,  $n=22$ , **Figure 1**). In the Cox model, combination  $\beta$ -glucan/antibody treatment was the only variable significantly associated with improved survival (hazard ratio treatment: 0.07, 95%CI 0.02, 0.27;  $p<0.0001$ ). Ten (45%) mice in the combination group survived long term with a median follow-up of 248 days. Only one mouse in any of the control groups (<5%) remained alive during the experiment. This anti-tumor effect was evident against a panel of GD2-positive neuroblastoma lines: NMB7, SK-N-JD, and SK-N-ER. Barley  $\beta$ -glucan was effective when the route of administration was either intragastric or intraperitoneal. In contrast, if the tumor was antigen-negative (e.g. GD2-negative rhabdomyosarcoma HTB82), 3F8 plus  $\beta$ -glucan treatment was ineffective. When the dose of oral  $\beta$ -glucan was decreased by 10-folds from 4000 ug to 400ug, 40 ug and 4 ug, the tumor growth rate were  $4.3 \pm 2.2\%$ ,  $3.8 \pm 0.9\%$ ,  $8.1 \pm 0.8\%$ , and  $9.5 \pm 0.9\%$ , respectively. These data suggest an optimal dose somewhere between 400 ug and 4000 ug. The animals did not suffer weight loss, or histopathologic changes in the major organs at necropsy in the treatment groups, irrespective of  $\beta$ -glucan dose. When the glucan was heated at 95°C up to 3 hours, its in vivo effects remained intact. However, following digestion with endo-(1 $\rightarrow$ 3), (1 $\rightarrow$ 4)- $\beta$ -D-glucanase, all its in vivo effect was destroyed. In addition, the anti-tumor effect of the antibodies was lost when the Fc of the antibody was removed by pepsin, or when an IgG1 isotype (Moab 8H9) was used.

Molecular size of (1→3), (1→4)- $\beta$ -D-glucan and anti-tumor synergy with MoAb. Barley  $\beta$ -glucans of increasing molecular sizes were tested at an oral dose of 40 ug or 400 ug. Anti-tumor effect improved with increasing molecular size, with a 2.2% decrease in tumor growth rate per day for each increase of 100kD in molecular weight; 95% confidence interval 3.0%, 1.4%;  $p < 0.00001$  (**Table 1**). However, since the shape of the  $\beta$ -glucan in aqueous solution correlated with average MW by MALLS analysis, potency could be a function of molecular shape rather than molecular size.

Source of  $\beta$ -D-glucan and anti-tumor synergy with MoAb. A (1→3), (1→4)- $\beta$ -D-glucan of average MW of 210 kD derived from barley was chosen as our standard. Using the neuroblastoma xenograft model, equivalent ug doses of  $\beta$ -glucans derived from various plant sources were compared in their anti-tumor activity when administered by intragastric injection plus intravenous MoAb 3F8 (**Table 1**). As expected, since the chemical composition (1→3), (1→4)- $\beta$ -D-glucan derived from barely, oat and wheat were similar, comparable levels of synergy with MoAb was found, and high MW also appeared to be more effective. When glucans with (1→6)- $\beta$ -linkages were tested, high MW species (e.g. Lentinan 1,500 kD) was not as effective compared to the standard. On the other hand, low MW (1→3), (1→6) preparations (e.g. Laminarin 5 kD), though not as effective as standard, was more potent than low MW (1→3), (1→4)- $\beta$ -D-glucan.

$\beta$ -glucan effect in a wide spectrum of MoAbs and human tumor models. Using the standard  $\beta$ -glucan from barley (210kD), a series of MoAb were screened against a panel of human tumor xenografts in various mouse strains. The combination of oral  $\beta$ -glucan with complement activating MoAb suppressed tumor growth significantly in contrast to MoAb or  $\beta$ -glucan alone. This was shown for anti-GD3 MoAb (R24) against

melanoma (**Figure 2A**), anti-EGF-R (528) MoAb against epidermoid carcinoma A431 (**Figure 2B**), and anti-HER2 (Herceptin) against human breast carcinoma BT474 xenografts in nude mice (**Figure 2C**). Again, MoAb 455, an IgG1 anti-EGF-R was ineffective against epidermoid carcinoma in contrast to the IgG2a 528 (**Figure 2B**). In metastatic lymphoma model, Daudi cells injected i.v. established widespread tumors in brain, spinal cord, kidneys and ovaries of SCID mice (data not shown). In the Cox model, only combination treatment and dose of Rituximab were associated with survival. Median survival was 59 days in animals receiving either Rituximab alone,  $\beta$ -glucan alone or no treatment. Median survival in the group treated with Rituximab plus  $\beta$ -glucan was 97 days (hazard ratio 0.09; 95% CI 0.03, 0.27;  $p < 0.001$ ).

### Discussion

We have shown that  $\beta$ -glucans greatly enhanced the anti-tumor effects of MoAb against established tumors in mice. We observed this effect irrespective of route of  $\beta$ -glucan administration (intragastric or intraperitoneal), antigen (GD2, GD3, CD20, EGFR, HER2), human tumor type (NB, melanoma, epidermoid carcinoma, lymphoma, breast cancer), mouse strain (athymic nu/nu, severe combined immune deficiency mice), or tumor site (subcutaneous versus systemic).  $\beta$ -glucan was heat-stable, its anti-tumor effect was dose- and schedule-dependent, requiring antibody-Fc, but not cytophilicity of the antibody. Neither antibody nor  $\beta$ -glucan alone was effective. We detected no toxicities even at  $\beta$ -glucan doses of 4000 ug/day for 4 weeks. This synergy of (1 $\rightarrow$ 3), (1 $\rightarrow$ 4)- $\beta$ -D-glucan with MoAb increased with  $\beta$ -glucan MW.

$\beta$ -glucans have been tested for tumor therapy in mice for nearly 40 years (24,25). Several forms of mushroom-derived

$\beta$ -glucans are used clinically in Japan to treat cancer, including polysaccharide Kureha (PSK, from *Coriolus versicolor*), Lentinan and Schizophyllan. In randomized trials in Japan, PSK and Schizophyllan have moderately  
5 improved survival rates in some cancer trials (26-30), and less encouraging in others (31,32). While  $\beta$ -glucans are not used by western oncologists,  $\beta$ -glucan containing botanical medicines such as Ling-Zhi, maitake and green barley are widely used by cancer patients in the US as  
10 alternative/complementary cancer therapies, often with poor clinical validation or quality control.

Given the biology of iC3b targeted cytotoxicity,  $\beta$ -glucan should have clinical potential. However, limitations with  
15 existing  $\beta$ -glucan strategies are several fold. They are generally expensive and inconvenient to administer: e.g. Lentinan and Schizophyllan are given i.v. daily over long periods of time. Besides being insoluble, they contain proteins and non- $\beta$ -glucan carbohydrates, which confound  
20 mechanistic studies and complicate the manufacturing and control process. Because of protein contaminants they are potentially allergenic. The spontaneous cross-linking of CR3 by  $\beta$ -glucan of high MW can cause neutrophil degranulation and cytokine release from macrophages,  
25 resulting in undesirable clinical toxicities. For low MW  $\beta$ -glucan, besides their low affinity for CR3, they have rapid renal clearance. Without anti-tumor antibodies to activate human complement,  $\beta$ -glucan is largely ineffective. Here we addressed these limitations by (1) using pure (1 $\rightarrow$ 3), (1 $\rightarrow$ 4)- $\beta$ -  
30 D-glucan from barley, (2) administering  $\beta$ -glucan orally instead of intravenously, and (3) coadministration of tumor-specific antibodies to ensure complement activation.

Previous studies have demonstrated that oral  $\beta$ -glucans  
35 activate splenic and peritoneal macrophages for tumor-

cytotoxicity. In a study of  $^{14}\text{C}$ -labeled oral  $\beta$ -glucan, sequestration in the liver was observed (33), suggesting that oral  $\beta$ -glucan entered the blood and behaved pharmacokinetically similar to intravenously administered low MW  $\beta$ -glucan (34-36). These studies also suggested that processing by the gastrointestinal tract produced  $\beta$ -glucan with high activity for CR3. Besides this model of intravasation of processed barley  $\beta$ -glucan, leukocytes could also be activated directly in the gut before homing to the tumor. It is of interest that unpurified  $\beta$ -glucan (Betratrim) has low biologic activity in our model. Despite the abundance of  $\beta$ -glucan (3% of the dry weight) in grains, its bioavailability from cereals is limited since high MW  $\beta$ -glucans requires high temperature ( $>60^\circ\text{C}$ ) extraction and final gelling. It is therefore not surprising that high-fiber (13.5 g/day) wheat-bran supplement did not have anti-tumor effects in recent human trials (37).

Our findings using (1 $\rightarrow$ 3), (1 $\rightarrow$ 4)- $\beta$ -D-glucans from barley were unexpected. In previous studies, the comb-like branch structure of (1 $\rightarrow$ 3), (1 $\rightarrow$ 6)- $\beta$ -linkage (e.g. lentinan, schizophyllan, laminarin, and  $\beta$ -glucan from Baker's yeast) was deemed requisite for its anti-tumor effect (38). In those models, however, T-cells were essential. In our studies, (1 $\rightarrow$ 3), (1 $\rightarrow$ 4)- $\beta$ -glucan could reproducibly enhance the anti-tumor effect of MoAb in immunodeficient mice, clearly demonstrating that neither T nor B cells were needed. Although the absolute proof of complement and CR3 requirement would have to await experiments with knock-out mice, preliminary evidence from studying MoAb isotypes and subclasses did suggest that complement activation was required. Since most cancers express mCRP (CD46, CD55, CD59) on their cell surface (39-46), complement mediated tumor lysis is typically inefficient. Nevertheless, despite these inhibitory proteins, iC3b has been detected on tumor

cells isolated from fresh human breast tumors, and enough levels could be deposited by MoAb (e.g. Herceptin) in vitro to opsonize tumor cells for phagocytes and NK cells in vitro (47). It is possible that sublethal levels of complement  
5 activation deposited enough iC3b to optimize tumor killing, a strategy that deserves clinical investigation.

## REFERENCES

1. Ross GD, Cain JA, Lachmann PJ. Membrane complement receptor type three (CR3) has lectin-like properties analogous to bovine conglutinin and functions as a receptor for zymosan and rabbit erythrocytes as well as a receptor for iC3b. J Immunol 1985;134:3307-15.
2. Ross GD, Cain JA, Myones BL, Newman SL, Lachmann PJ. Specificity of membrane complement receptor type three (CR3) for beta-glucans. Complement Inflamm 1987;4:61-74.
3. Cain JA, Newman SL, Ross GD. Role of complement receptor type three and serum opsonins in the neutrophil response to yeast. Complement Inflamm 1987;4:75-86.
4. Vetvicka V, Thornton BP, Wieman TJ, Ross GD. Targeting of natural killer cells to mammary carcinoma via naturally occurring tumor cell-bound iC3b and beta-glucan-primed CR3 (CD11b/CD18). J Immunol 1997;159:599-605.
5. Irie K, Irie RF, Morton DL. Evidence for in vivo reaction of antibody and complement to surface antigens of human cancer cells. Science 1974;186:454-6.
6. Seegal BC, Hsu KC, Lattimer JK, Habif DV, Tannenbaum M. Immunoglobulins, complement and foreign antigens in human tumor cells. Int Arch Allergy Immunol 1976;52:205-11.
7. McCoy JP, Hofheinz DE, Ng AB, Nordqvist S, Haines HG. Tumor-bound immunoglobulin in human gynecologic cancers. J. Natl. Cancer Inst. 1979;63:279-83.

8. Thornton BP, Vetvicka V, Pitman M, Goldman RC, Ross GD. Analysis of the sugar specificity and molecular location of the B-glucan-binding lectin site of complement receptor type 3 (CD11b/CD18). J Immunol 1996;156:1235-46.
9. Xia Y, Vetvicka V, Yan J, al. e. The beta-glucan-binding lectin site of mouse CR3 (CD11b/CD18) and its function in generating a primed state of the receptor that mediates cytotoxic activation in response in iC3b-opsonized target cells. J Immunol 1999;162:2281-90.
10. Vetvicka V, Thornton BP, Ross GD. Soluble beta-glucan polysaccharide binding to the lectin site of neutrophil or natural killer cell complement receptor type 3 (CD11b/CD18) generates a primed state of the receptor capable of mediating cytotoxicity of iC3b-opsonized target cells. Journal of Clinical Investigations 1996;98:50-61.
11. Yan J, Vetvicka V, Xia Y, Coxon A, Carroll MC, Mayadas TN, et al. B-glucan a "Specific" biologic response modifier that uses antibodies to target tumors for cytotoxic recognition by leukocyte complement receptor type 3 (CD11b/CD18). J Immunol 1999;163:3045-52.
12. Di Renzo L, Yefenof E, Klein E. The function of human NK cells is enhanced by beta-glucan, a ligand of CR3 (CD11b/CD18). Eur J Clin Nutr 1991;21:1755-8.
13. Czop JK, Austen KF. Properties of glycans that activate the human alternative complement pathway and interact with the human monocyte beta-glucan receptor. J Immunol 1985;135:3388-93.
14. Thornton BP, Vetvicka V, Ross GD. Function of C3 in a humoral response: iC3b/C3dg bound to an immune complex



- generated with natural antibody and a primary antigen promotes antigen uptake and the expression of co-stimulatory molecules by all B cells, but only stimulates immunoglobulin synthesis by antigen-specific B cells. Clinical Experimental Immunology 1996;104:531-7.
15. Roubroeks JP, Skjak-Braek G, Ryan L, Christensen BE. Molecular weight dependency on the production of the TNF stimulated by fractions of rye (1-->3), (1-->4)-beta-D-glucan. Scand J Immunol 2000;52:584-7.
16. Cheung NK, Saarinen U, Neely J, Landmeier B, Donovan D, Coccia P. Monoclonal antibodies to a glycolipid antigen on human neuroblastoma cells. Cancer Research 1985;45:2642-9.
17. Modak S, Kramer K, Humayun G, Guo HF, Cheung NKV. Monoclonal antibody 8H9 targets a novel cell surface antigen expressed by a wide spectrum of human solid tumors. Cancer Research 2001;61:4048-54.
18. Houghton AN, Mintzer D, Cordon-Cardo C, Welt S, Fliegel B, Vadhsans S, et al. Mouse monoclonal antibody detecting GD3 ganglioside: a phase I trial in patients with malignant melanoma. Proc. Natl Acad Sci, USA 1985;82:1242-6.
19. Masui H, Moroyama T, Mendelsohn J. Mechanism of antitumor activity in mice for anti-epidermal growth factor receptor monoclonal antibodies with different isotypes. Cancer Res 1986;46:5592-8.
20. Ciucanu I, Kerek F. A simple and rapid method for the permethylation of carbohydrates. Carbohydrate Res 1984;131:209-17.

21. Wyatt P. Light scattering and the absolute characterization of macromolecules. *Analytical Chimica Acta* 1993;272:1-40.
- 5 22. Knuckles BE, Yokohama WH, Chiu MM. Molecular characterization of barley  $\beta$ -glucans by size-exclusion chromatography with multiple-angle laser-light scattering and other detectors. *Cereal Chemistry* 1997;74:599-604.
- 10 23. Bergman I, Basse PH, Barmada MA, Griffin JA, Cheung NKV. Comparison of in vitro antibody-targeted cytotoxicity using mouse, rat and human effectors. *Cancer Immunol. Immunother.* 2000;49:259-66.
- 15 24. Diller IC, Mankowski ZT, Fisher ME. The effect of yeast polysaccharides on mouse tumors. *Cancer Res* 1963;23:201-8.
- 20 25. Sveinbjornsson B, Rushfeldt C, Seljelid R, Smedsrod B. Inhibition of establishment and growth of mouse liver metastases after treatment with interferon gamma and beta-1,3-D-Glucan. *Hepatology* 1998;27(5):1241-8.
- 25 26. Niimoto M, Hattori T, Tamada R, Sugimachi K, Inokuchi K, Ogawa N. Postoperative adjuvant immunochemotherapy with mitomycin C, fluorouracil, and PSK for gastric cancer. An analysis of data on 579 patients followed for five years. *Japanese Journal of Surgery* 1988;18:681-6.
- 30 27. Mitomi T, Tsuchiya S, Iijima N, Aso K, Suzuki K, Nishiyama K, et al. Randomized, controlled study on adjuvant immunochemotherapy with PSK in curatively resected colorectal cancer. The cooperative study group of surgical adjuvant immunochemotherapy for cancer of colon and rectum (Kanagawa). *Dis Colon Rectum* 1992;35:123-30.
- 35

28. Ogoshi K, Satou H, Isono K, Mitomi T, Endoh M, Sugita M. Immunotherapy for esophageal cancer. A randomized trial in combination with radiotherapy and radiochemotherapy. Cooperative study group for esophageal cancer in Japan. American Journal of Clinical Oncology 1995;18:216-22.
29. Fujimoto S, Furue H, Kimura T, Kondo T, Orita K, Taguchi T, et al. Clinical outcome of postoperative adjuvant immunochemotherapy with sizofiran for patients with resectable gastric cancer: a randomised controlled study. Eur J Cancer 1991;27:1114-8.
30. Okamura K, Suzuki M, Chihara T, Fujiwara A, Fukada T, Goto S, et al. Clinical evaluation of sizofiran combined with irradiation in patients with cervical cancer. A randomized controlled study; a five-year survival rate. Biotherapy 1989;1:103-7.
31. Toi M, Hattori T, Akagi M, Inokuchi K, Orita K, Sugimachi K, et al. Randomized adjuvant trial to evaluate the addition of tamoxifen and PSK to chemotherapy in patients with primary breast cancer. 5-year results from the Nishi-Nippon group of the adjuvant chemoendocrine therapy for breast cancer organization. Cancer 1992;70:2475-83.
32. Ohno R, Yamada K, Masaoka T, al. e. A randomized trial of chemoimmunotherapy of acute nonlymphocytic leukemia in adults using a protein-bound polysaccharide preparation. Cancer Immunol. Immunother. 1984;18:149-54.
33. Ikuzawa M, Matsunaga K, Nishiyama S, Nakajima S, Kobayashi Y, Andoh T, et al. Fate and distribution of an

antitumor protein-bound polysaccharide PSK (Krestin).  
Int. J. Immunopharmacol. 1988;10:415-23.

34. Yan J, Vetvicka V, Xia Y, Hanikyrova M, Mayadas TN,  
5 Ross GD. Critical role of kupffer cell CR3 (CD11b/CD18)  
in the clearance of IgM-opsonized erythrocytes or  
soluble beta-glucan. Immunopharmacology 2000;46:39-54.
35. Smedsrod B, Seljelid R. Fate of intravenously injected  
10 aminated beta(1->3) polyglucose derivatized with 125I-  
tyraminyl cellobiose. Immunopharmacology 1991;21:149-  
58.
36. Suda M, Ohno N, Adachi Y, Yadomae T. Relationship  
15 between the tissue distribution and antitumor activity  
of highly branched (1-->3)-beta-D-glucan, SSG. Biol.  
Pharm. Bull. 1994;17:131-5.
37. Alberts DSea. Lack of effects of a high-fiber cereal  
20 supplement on the recurrence of colorectal adenomas. N  
Engl J Med 2000;342:1156-62.
38. Bohn JA, BeMiller JN. (1-3)-B-D-Glucans as biological  
response modifiers:a review of structure-functional  
25 activity relationships. Carbohydr Polymers 1995;28:3-  
14.
39. Koretz K, Bruderlein S, Henne C, Moller P. Expression  
of CD59, a complement regulator protein and a second  
30 ligand of the CD2 molecule, and CD46 in normal and  
neoplastic colorectal epithelium. Br J Cancer  
1993;68:926-31.
40. Juhl H, Helmig F, Baltzer K, Kalthoff H, Henne-Bruns D,  
35 Kremer B. Frequent Expression of Complement Resistance  
Factors CD46,CD55,and CD55 on Gastrointestinal Cancer  
Cells Limits the Therapeutic Potential of Monoclonal

Antibody 17-1A. Journals of Surgical Oncology  
1997;64:222-30.

41. Brasoveanu LI, Altomonte M, Gloghini A, Fonsatti E,  
5 Coral S, Gasparollo A, et al. Expression of protectib  
(CD59) in human melanoma and its functional role in  
cell-and complement-cytotoxicity. Int J Cancer  
1995;61:548-56.
- 10 42. Hara T, Kojiama A, Fukuda H, Masaoka T, Fukumori Y,  
Matsumoto M, et al. Levels of complement regulator  
protein,CD35(CR1),CD46(MCP) and CD55(DAF) in human  
haematological malignancies. British Journal of  
Haemotolgy 1992;82:368-73.
- 15 43. Marquart HV, Gronbaek K, Christensen BE, Svehag S-E,  
Lesli RGQ. Complement activation by malignant B-cells  
from patients with chronic lymphocytic leukemia.  
Clinical Experimental Immunology 1995;102:575-81.
- 20 44. Bjorge L, Hakulinen J, Wahlstrom T, Matre R, Meri S.  
Complement-regulatory proteins on ovarian malignancies.  
Int J Cancer 1997;70:14-25.
- 25 45. Jurianz K, Maslak S, Garcia-Schuler H, Fishelson Z,  
Kirschfink M. Neutralization of complement regulatory  
proteins augments lysis of breast carcinoma cells  
targeted with rhumAb anti-HER2. Immunopharmacology  
1999;42:209-18.
- 30 46. Jurianz K, Ziegler S, Garcia-Schuler H, Kraus S,  
Bohana-Kashtan O, Fishelson Z, et al. Complement  
resistance of tumor cells: basal and induced  
mechanisms. Mol Immunol 1999;36:929-39.

47. Klein E, DiRenzo L, Yefenof E. Contribution of C3,CD11b/CD18 to cytolysis by human NK cells. Mol Immunol 1990;27:1343-7.

**Table 1:** Neuroblastoma growth rate (%/day) when treated with i.v. 3F8 and oral  $\beta$ -glucan derived from various plant sources.

5			<u>400 ug****</u> <u>40 ug****</u>	
	<u>Glucans</u>	<u>Description</u>	<u>MW (kD)</u>	<u>% tumor growth relative to standard</u>
	<b>(1<math>\rightarrow</math>3), (1<math>\rightarrow</math>4) -<math>\beta</math>-D-glucan*</b>			
	Standard barley glucan		210	100
	Antibody alone Control		-	287
10	Saline control		-	307
	Barley	MW standard	337	79
		MW standard	237	100
		MW standard	178	117
		MW standard	131	163
15		MW standard	48	180
		BBG111	266	60
		BBG126	90	146
	Oat	BBG128	262	88
		BBG127	201	104
20	Wheat	BBG117	138	-
	Betatrim	Unpurified	-	325
	Lichenan	BBG113	132	189
	<b>(163), (166) -<math>\beta</math>-D-glucan</b>			
	Laminarin**	BBG108	5	177
25	Laminarin***	BBG109	32	326
	Lentinan	BBG114	1491	123

\*(163) $\beta$ - linkage was ~30% for all the (163), (164)- $\beta$ -D-glucans

\*\* (163) $\beta$ -linkage was 92%

30 \*\*\* (163) $\beta$ -linkage was 53%

\*\*\*\* Either 400 or 40 ug of (163)- $\beta$ -D-glucans was administered orally qd x 29 days plus intravenous MoAb 3F8 twice a week (M,Th) x 8 doses in groups of 4-5 mice each. Tumor size was measured periodically over the entire treatment period. Tumor growth curve and potency were calculated as detailed in Materials and Methods.

35

**What is claimed is:**

1. A composition comprising an effective amount of glucan capable of enhancing efficacy of antibodies.
- 5 2. The composition of claim 1, wherein the antibody is a monoclonal antibody.
3. The composition of claim 1, wherein the antibody is an antibody against cancer.
- 10 4. The composition of claim 3, wherein the antibody is a tumor-binding antibody.
- 15 5. The composition of claim 4, wherein the antibody is capable of activating complement.
6. The composition of claim 5, the antibody is further capable of activating the antibody dependent cell-mediated cytotoxicity.
- 20 7. The composition of claim 4, wherein the antibody is directed at the epidermal growth factor receptor.
- 25 8. The composition of claim 7, wherein the antibody is 528 or C225.
9. The composition of claim 4, wherein the antibody is directed to a ganglioside.
- 30 10. The composition of claim 9, wherein the ganglioside is GD3.
11. The composition of claim 10, wherein the antibody is R24.
- 35 12. The composition of claim 4, wherein the ganglioside is GD2.



13. The composition of claim 12, wherein antibody is 3F8.
14. The composition of claim 4, wherein the antigen is CD20.
- 5 15. The composition of claim 14, wherein the antibody is Rituximab.
- 10 16. The composition of claim 4, wherein the antigen is CD25.
17. The composition of claim 16, wherein the antibody is Dacluzimab.
- 15 18. The composition of claim 4, wherein the antigen is Her2/neu.
19. The composition of claim 18, wherein the antibody is Herceptin.
- 20 20. The composition of claim 4, wherein the antigen is CD22.
21. The composition of claim 20, wherein the antibody is Epratuzumab.
- 25 22. The composition of claim 3, wherein the cancer is neuroblastoma, melanoma, non-Hodgkin's lymphoma, Epstein-Barr related lymphoma, Hodgkin's lymphoma, retinoblastoma, small cell lung cancer, brain tumors, leukemia, epidermoid carcinoma, prostate cancer, renal cell carcinoma, transitional cell carcinoma, breast cancer, ovarian cancer, lung cancer, colon cancer, liver cancer, stomach cancer, or other gastrointestinal cancers.
- 30 23. The composition of claim 1 and a pharmaceutically acceptable carrier.
- 35

24. A method for treating a subject comprising administering the composition of claim 1-23 to the subject.
- 5 25. A composition comprising effective amount of glucan capable of enhancing efficacy of vaccines.
26. A composition of claim 25, wherein the vaccine is  
10 against cancer.
27. The composition of claim 25 and a pharmaceutically acceptable carrier.
- 15 28. A method of treating a subject comprising administering the composition of claim 25 to the subject.
29. A composition of claim 25, wherein the vaccine is  
20 against infectious agents.
30. A composition of claim 29, wherein the infectious agents are bacteria, viruses, fungi, or parasites.
- 25 31. A composition comprising effective amount of glucan capable of enhancing efficacy of natural antibodies.
32. The composition of claim 31, wherein the antibodies are  
30 against cancer.
33. The composition of claim 31, wherein the antibodies are  
35 against infectious agents.
34. The composition of claim 31, wherein the infectious  
agents are bacteria, viruses, fungi, or parasites.
35. A composition comprising effective amount of glucan capable of enhancing host immunity.

36. The composition of claim 35, wherein the immunity is against cancer.
- 5 37. The composition of claim 35, wherein the immunity is against infectious agents.
38. A composition comprising effective amount of glucan capable of enhancing the action of an agent in preventing tissue rejection.
- 10 39. The composition of claim 38, wherein the agent is an antibody.
- 15 40. The composition of claim 39, wherein the antibody modulates T-cell function.
41. The composition of claim 40, wherein the antibody is anti-CD25 or anti-CD3.
- 20 42. The composition of claim 39, wherein the antibody modulates B-cell function.
43. The composition of claim 42, wherein the antibody is anti-CD20.
- 25 44. The composition of claim 38, wherein the tissue is transplanted tissue or transplanted organ.
- 30 45. The composition of claim 38, wherein the tissue is the host as in graft-versus-host disease.
46. The composition of claim 1, 25, 31, 35, or 38 wherein the glucan are. 1,3-1,4 mixed linkage, without 1,6 branches.
- 35 47. The composition of claim 1, 25, 31, 35, or 38 wherein the glucan is of high molecular weight.

48. The composition of claim 47, wherein the molecular weight of the glucan ranges from 250,000 to 450,000 daltons.
- 5 49. The composition of claim 1, 25, 31, 35, or 38 wherein the glucan is derived from barley, oat, wheat or moss.
50. The composition of claim 1, 25, 31, 35, or 38 wherein the glucan is stable to heat treatment.
- 10 51. The composition of claim 50, wherein the composition is stable after boiling for 3 hours.
52. The composition of claim 1, 25, 31, 35, or 38 when oral  
15 route is adopted when taken into a subject.
53. The composition of claim 52, wherein the effective dose is about  $\geq 25$  mg/kg/day, five days a week for a total of 2-4 weeks.
- 20 54. A method for treating a subject comprising administering the composition of claims 1-53 to the subject.

Figure 1




Figure 2




Figure 3





Figure 4





5/30

Figure 5




Figure 6




Figure 7




Figure 8




Figure 9




Figure 10





Figure 11



12/30

Figure 12






Figure 13




Figure 14




Figure 15




Figure 16




Figure 17




Figure 18




Figure: 19




Figure 20






Figure 1A

glucan-nb.doc






Figure 1B




glucan-nb.doc

Figure 1C




glucan-nb.doc

Figure 2



glucan-nb.doc

Figure 3



26/30

glucan-nb.doc

Figure 4




Figure 5

glucan-nb.doc





Figure 6


glucan-nb.doc






29/30

Figure 1



30/30

Figure 2



## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US02/01276

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : A61K 31/715

US CL : 514/54

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/54

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CAPLUS, USPATFUL, PCTFUL, MEDLINE

search terms: glucan, antibody, cancer, vaccine, immunity, complement, growth factor, etc.

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X --- A	US 5,189,028 A (NIKL et al) 23 February 1993, see entire text, especially columns 1- 2.	25-37 ----- 1-24, 38-54
X ---- A	US 4,833,131 A (WILLIAMS et al) 23 May 1989, see entire text, especially abstract and columns 8-10.	1-3, 6, 22-24, 35,46-54 ----- 4-5, 7-21, 25-34, 36-45

☐ Further documents are listed in the continuation of Box C.
 ☐ See patent family annex.

* Special categories of cited documents:	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A" document defining the general state of the art which is not considered to be of particular relevance	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"E" earlier document published on or after the international filing date	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"&" document member of the same patent family
"O" document referring to an oral disclosure, use, exhibition or other means	
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

16 MAY 2002

Date of mailing of the international search report

05 JUN 2002

 Name and mailing address of the ISA/US  
 Commissioner of Patents and Trademarks  
 Box PCT  
 Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

VICKIE KIM

Telephone No. (703) 3081234